

IDSA/PIDS announce guidelines for treating pneumonia in children

August 31 2011

Immunizations, including a yearly flu vaccine, are the best way to protect children from life-threatening pneumonia, according to new guidelines from the Pediatric Infectious Diseases Society (PIDS) and the Infectious Diseases Society of America (IDSA).

The guidelines, which are the first on diagnosing and treating community-acquired pneumonia (CAP) in infants and [children](#), place preventing bacterial pneumonia as a top priority.

Every year, pneumonia kills more than 2 million children ages 5 years and younger worldwide. In the United States, 525 children 15 years old or younger died from pneumonia and other lower respiratory infections in 2006, according to the [Centers for Disease Control and Prevention](#) (CDC). While pneumonia can be caused by different types of bugs, in infants and preschool children it usually is caused by a virus, which doesn't need to be treated with antibiotics. However, antibiotics are needed for bacterial pneumonia, which is the most serious type.

Although there are guidelines for diagnosing and treating pneumonia in adults, the course of bacterial pneumonia tends to be different for children. Because of this, practices vary from hospital to hospital, and doctor to doctor. The guidelines from PIDS and IDSA provide all physicians who care for children with a roadmap to the most scientifically valid diagnosis and treatment recommendations.

"Diagnostic methods and treatments that work well in adults may be too

risky and not have the desired result in children," said John S. Bradley, MD, lead author of the CAP guidelines and professor and chief of the division of [infectious diseases](#) at the University of California at San Diego Department of Pediatrics. "With these guidelines, we are hopeful that the standard and quality of care children receive for community-acquired pneumonia will be consistent from doctor to doctor – providing much better treatment outcomes."

Because viral infections such as influenza can develop into bacterial pneumonia, it's important that children 6 months and older receive a yearly influenza vaccine, according to the guidelines. It is also important that infants and children are up to date on their other scheduled vaccines, several of which prevent bacterial pneumonia. The successful U.S. vaccination program has significantly reduced bacterial pneumonia, and therefore has prevented deaths from the infection, notes Dr. Bradley.

While the guidelines stress the importance of diagnosing pneumonia appropriately, they also warn that over-treatment is a critical concern. For instance, most pneumonia in preschool-aged children is viral, meaning it will run its course and will not develop into life-threatening bacterial pneumonia. In these cases, there is no need to perform unnecessary medical interventions such as using x-rays (which expose the child to radiation needlessly) or prescribing antibiotics (which kill bacteria, not viruses, and may foster drug-resistant bacteria).

"A child with chest congestion, a cough, runny nose and low-grade fever likely has viral pneumonia, and Mother Nature treats those herself," said Dr. Bradley. "If the child has a fever of 104, is barely able to keep fluids down, just wants to lie in bed and is breathing fast, it may be bacterial pneumonia and require antibiotics and hospitalization."

The guidelines suggest when doctors can feel comfortable not prescribing a higher level of care and when they need to be cautious, and

do so. By design, the guidelines lead off with recommendations regarding hospitalization. "Most of these kids will have their first encounter when they have fever and difficulty breathing and see their primary care physician, or the emergency room doctor," said Dr. Bradley. "The first major decision that needs to be made is, is this child well enough to go home, or does he or she need a higher level of care?"

For instance, the guidelines recommend infants 3 to 6 months old with suspected bacterial pneumonia are likely to benefit from hospitalization, even if the pneumonia isn't confirmed by blood tests. Blood testing in children often isn't accurate, so physicians need to pay close attention to symptoms, and, if unsure, err on the side of treating, said Dr. Bradley.

Following are some other recommendations included in the guidelines:

- Because infants 6 months and younger cannot get the flu shot or nasal spray, their parents and caregivers should be sure to get the vaccine.
- When antibiotics are necessary, amoxicillin should be first-line therapy for [bacterial pneumonia](#), because it is safe and effective. Many doctors prescribe more powerful antibiotics, which are unnecessary and can kill off good bacteria in the body.
- Although pneumonia from methicillin-resistant *Staphylococcus aureus* (MRSA) is uncommon, it can cause severe illness, so physicians need to consider it if a child doesn't improve after first-line antibiotic therapy.

For each of the 92 specific recommendations, the guidelines denote the strength of the recommendation as well as the quality of evidence for each. The guidelines note the lack of solid evidence in some areas – often due to the ethical challenges of studying children – and call for research in specific areas.

"We're hopeful that in following these guidelines, physicians and hospitals will collect data and the results can be compared," said Dr. Bradley. "We envision this as the first of many revisions of guidelines to come."

The 13-member guidelines panel was comprised of experts from around the country, including lung, emergency department, hospital medicine and critical care specialists, office-based pediatricians, pediatric surgeons and CDC epidemiologists. They reviewed hundreds of scientific studies, papers and presentations in preparation for writing the [guidelines](#). In addition to Dr. Bradley, the panel included: Carrie L. Byington, Samir S. Shah, Brian Alverson, Edward R. Carter, Christopher Harrison, Sheldon L. Kaplan, Sharon E. Mace, George H. McCracken, Jr., Matthew R. Moore, Shawn D. St. Peter, Jana A. Stockwell and Jack T. Swanson.

Provided by Infectious Diseases Society of America

Citation: IDSA/PIDS announce guidelines for treating pneumonia in children (2011, August 31) retrieved 26 April 2024 from <https://medicalxpress.com/news/2011-08-idsapids-guidelines-pneumonia-children.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--