

Novel imaging probe allows noninvasive detection of dangerous heart-valve infection

August 21 2011

A novel imaging probe developed by Massachusetts General Hospital (MGH) investigators may make it possible to diagnose accurately a dangerous infection of the heart valves. In their *Nature Medicine* report, which is receiving advance online publication, the team from the MGH Center for Systems Biology describes how the presence of *Staphylococcus aureus*-associated endocarditis in a mouse model was revealed by PET imaging with a radiolabeled version of a protein involved in a process that usually conceals infecting bacteria from the immune system.

"Our probe was able to sense whether *S. aureus* was present in abnormal growths that hinder the normal function of heart valves," says Matthias Nahrendorf, MD, PhD, of the MGH Center for Systems Biology, a co-lead author of the study. "It has been very difficult to identify the bacteria involved in endocarditis, but a precise diagnosis is important to steering well-adjusted <u>antibiotic therapy</u>."

An infection of the tissue lining the heart valves, endocarditis is characterized by growths called vegetations made up of clotting components such as platelets and fibrin along with infecting microorganisms. Endocarditis caused by *S. aureus* is the most dangerous, with a mortality rate of from 25 to almost 50 percent, but diagnosis can be difficult since symptoms such as fever and heart murmur are vague and blood tests may not detect the involved bacteria. Without appropriate antibiotic therapy, *S. aureus* endocarditis can progress rapidly, damaging or destroying <u>heart valves</u>.



S. aureus bacteria initiate the growth of vegetations by secreting staphylocoagulase, an enzyme that sets off the clotting cascade. This process involves a protein called prothrombin, which is part of a pathway leading to the deposition of fibrin, a primary component of blood clots. The clotting process enlarges the vegetation, anchors it to the heart valve and serves to conceal the bacteria from <u>immune cells</u> in the bloodstream.

To develop an imaging-based approach to diagnosing *S. aureus* endocarditis, the MGH team first investigated the molecular mechanism by which staphylocoagulase sets off the clotting cascade, finding that one staphylocoagulase molecule interacts with at least four molecules of fibrin or its predecessor molecule fibrinogen in a complex that binds to a growing vegetation. Since prothrombin is an essential intermediary in the staphylocoagulase/fibrin interaction, the researchers investigated whether labeled versions of prothrombin could accurately detect *S. aureus* endocarditis in mice.

After initial experiments confirmed that an optical imaging technology called FMT-CT could detect a fluorescence-labeled version of prothombin deposited into *S. aureus*-induced vegetations, the researchers showed that a radiolabeled version of prothombin enabled the detection of *S. aureus* vegetations with combined PET-CT imaging, an approach that could be used in human patients after additional development and FDA approval.

"An approach like this could help clinicians detect the presence of endocarditis, determine its severity and whether it is caused by *S. aureus*, and track the effectiveness of antibiotics or other treatments," says Nahrendorf, also a co-corresponding author of the <u>Nature Medicine</u> article and an assistant professor of Radiology at Harvard Medical School. "We are working to improve the PET reporter probe with streamlined chemistry and a more mainstream PET isotope to make it a better candidate for eventual testing in patients."



Provided by Massachusetts General Hospital

Citation: Novel imaging probe allows noninvasive detection of dangerous heart-valve infection (2011, August 21) retrieved 26 April 2024 from https://medicalxpress.com/news/2011-08-imaging-probe-noninvasive-dangerous-heart-valve.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.