

# Poultry farms that go organic have significantly fewer antibiotic-resistant bacteria

August 10 2011

---



In this study, poultry houses like this, that transitioned to organic practices and ceased using antibiotics, had significantly lower levels of drug-resistant and multi-drug resistant enterococci bacteria. Credit: Amy R. Sapkota, University of Maryland


(Medical Xpress) -- Antibiotic use in conventional animal food production in the United States has created public health concern because it has been shown to contribute to the development of antibiotic-resistant bacteria that can potentially spread to humans. A new study, led by Dr. Amy R. Sapkota of the University of Maryland School of Public

Health, provides data demonstrating that poultry farms that have transitioned from conventional to organic practices and ceased using antibiotics have significantly lower levels of drug-resistant enterococci bacteria.

The study, published in [Environmental Health Perspectives](#) (online August 10, 2011), is the first to demonstrate lower levels of [drug-resistant bacteria](#) on newly organic farms in the United States and suggests that removing antibiotic use from large-scale U.S. [poultry farms](#) can result in immediate and significant reductions in [antibiotic resistance](#) for some bacteria.

"We initially hypothesized that we would see some differences in on-farm levels of antibiotic-resistant enterococci when poultry farms transitioned to organic practices. But we were surprised to see that the differences were so significant across several different classes of antibiotics even in the very first flock that was produced after the transition to organic standards," explains Sapkota, an Assistant Professor with the Maryland Institute for Applied Environmental Health. "It is very encouraging."

Sapkota and her team, which included R. Michael Hulet (Pennsylvania State University), Guangyu Zhang, Sam Joseph and Erinna Kinney (University of Maryland), and Kellogg J. Schwab (Johns Hopkins Bloomberg School of [Public Health](#)), investigated the impact of removing antibiotics from U.S. poultry farms by studying ten conventional and ten newly organic large-scale poultry houses in the mid-Atlantic region. They tested for the presence of enterococci bacteria in poultry litter, feed, and water, and tested its resistance to 17 common antimicrobials.



Multi-drug resistant bacteria are of particular public health concern because they can be resistant to all available antibiotics, and are therefore very difficult to treat if contracted by an animal or human. In this study, 42 percent of *Enterococcus faecalis* from conventional farms were multi-drug resistant, compared to only 10 percent from newly organic farms, and 84 percent of *Enterococcus faecium* from conventional farms were multi-drug resistant compared to 17 percent of those from newly organic farms. Credit: Amy R. Sapkota, University of Maryland

"We chose to study enterococci because these microorganisms are found in all poultry, including poultry on both organic and conventional farms. The enterococci are also notable opportunistic pathogens in human patients staying in hospitals. In addition, many of the antibiotics given in feed to farm animals are active against Gram-positive bacteria such as the enterococci. These features, along with their reputation of easily exchanging resistance genes with other bacteria, make enterococci a good model for studying the impact of changes in [antibiotic use](#) on farms," Sapkota explains.

While all farms tested positive for the presence of enterococci in [poultry litter](#), feed, and water as expected, the newly organic farms were characterized by a significantly lower prevalence of antibiotic-resistant

enterococci. For example, 67% of *Enterococcus faecalis* recovered from conventional poultry farms were resistant to erythromycin, while 18% of *Enterococcus faecalis* from newly organic poultry farms were resistant to this antibiotic. Dramatic changes also were observed in the levels of multi-drug [resistant bacteria](#) (organisms resistant to three or more antimicrobial classes) on the newly organic farms. Forty-two percent of *Enterococcus faecalis* from conventional farms were multi-drug resistant, compared to only 10% from newly organic farms, and 84% of *Enterococcus faecium* from conventional farms were multi-drug resistant compared to 17% of those from newly [organic farms](#) [see figure]. Multi-drug resistant bacteria are of particular public health concern because they can be resistant to all available antibiotics, and are therefore very difficult to treat if contracted by an animal or human.

"While we know that the dynamics of antibiotic resistance differ by bacterium and antibiotic, these findings show that, at least in the case of enterococci, we begin to reverse resistance on farms even among the first group of animals that are grown without antibiotics. Now we need to look forward and see what happens over 5 years, 10 years in time," says Sapkota.

She expects that reductions in drug-resistant bacteria on US farms that "go organic" are likely to be more dramatic over time as reservoirs of resistant [bacteria](#) in the farm environment diminish.

Provided by University of Maryland

Citation: Poultry farms that go organic have significantly fewer antibiotic-resistant bacteria (2011, August 10) retrieved 10 April 2024 from <https://medicalxpress.com/news/2011-08-poultry-farms-significantly-antibiotic-resistant-bacteria.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.