


Brain waves control the impact of noise on sleep

September 6 2011



Left: Auditory brain regions remain active in response to sounds during human non-rapid eye movement sleep. Right: When sounds occur during brain waves called sleep 'spindles', their transmission to the auditory cortex is distorted. Only a small region in the brainstem (arrow) remains activated in response to sounds. (c) ULg CRC

During sleep, our perception of the environment decreases. However the extent to which the human brain responds to surrounding noises during sleep remains unclear. In a study published this week in *Proceedings of the National Academy of Sciences (PNAS)*, researchers from University of Liege (Belgium) used brain imaging to study responses to sounds during sleep. They show that brain activity in the face of noise is

controlled by specific brain waves during sleep. In particular, waves called sleep ‘spindles’ prevent the transmission of sounds to auditory brain regions. Conversely, when sounds are associated with brain waves called ‘K-complexes’, activation of auditory areas is larger. Our perception of the environment is therefore not continuously reduced during sleep, but rather varies throughout sleep under the influence of particular brain waves.

In this study, the research team led by Dr Thanh Dang-Vu and Prof. Pierre Maquet (Cyclotron Research Center, University of Liège) shows that [brain activity](#) induced by sounds during sleep closely depends on brain waves that constitute our sleep.

By using functional magnetic resonance imaging (fMRI) combined with electroencephalography (EEG), researchers have evidenced that auditory brain regions remain active in response to sounds during sleep [see image, left panels], except when sounds occur during brain waves called sleep ‘spindles’. The study indeed shows that spindles prevent the transmission of sounds to the auditory cortex [see image, right panels].

Conversely, sounds can induce the production during sleep of brain waves called ‘K-complexes’. The results brought by this new study demonstrate that production of K-complexes by sounds is associated with a larger activation of auditory brain areas. While spindles prevent the transmission of sounds, K-complexes reflect a more important transmission of sounds to the sleeping brain.

The effects of noise on sleep are therefore controlled by specific brain waves. In particular, the human brain is isolated from the environment during sleep spindles, which might allow essential sleep functions to operate such as the consolidation of memory for previously acquired information. These [brain waves](#) thus play a crucial role in sleep quality and stability in the face of noise.

More information: “Interplay between spontaneous and induced brain activity during human non-rapid eye movement sleep”, *PNAS*, 2011, doi 10.1073, by T.T. Dang-Vu, M. Bonjean, M. Schabus, M. Boly, A. Darsaud, M. Desseilles, C. Degueldre, E. Balteau, C. Phillips, A. Luxen, T. Sejnowski et P. Maquet.

Provided by University of Liege

Citation: Brain waves control the impact of noise on sleep (2011, September 6) retrieved 2 May 2024 from <https://medicalxpress.com/news/2011-09-brain-impact-noise.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--