

# Lower contrast agent dose feasible in 320 row CT angiography

October 1 2011

---

The analysis of 180 CT angiography studies done using a 320 detector row CT scanner found that a contrast media protocol based on 60 milliliters of iopamidol "had sufficient enhancement in more than 96% of coronary segments," said Frank Rybicki, MD, of Brigham and Women's Hospital in Boston, and one of the authors of the study.

"Many centers currently use a higher iodine load comparable or equal to 80 milliliters of iopamidol," he said. The study shows that "this is not necessary, and the extra contrast means unnecessary costs and increased risk of contrast-induced nephropathy (CIN) to the patients." The risk of CIN is of special concern for these patients because they commonly have a renal insufficiency, which is known major risk factor for CIN, Dr. Rybicki added.

"We analyzed larger patients only (those with a [body mass index](#) of more than 30) and found that 92% of coronary segments had sufficient enhancement," said Dr. Rybicki. "These results support the general use of 60 milliliters of iopamidol for [CT angiography](#) done on 320 detector row CT scanners," he said.

The study is published in the October, 2011 [American Journal of Roentgenology](#).

Provided by American Roentgen Ray Society

Citation: Lower contrast agent dose feasible in 320 row CT angiography (2011, October 1)  
retrieved 27 April 2024 from

<https://medicalxpress.com/news/2011-10-contrast-agent-dose-feasible-row.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.