

Cough may warn of danger for patients with lung-scarring disease

October 18 2011

A new analysis has found that coughing may signal trouble for patients with the lung-scarring disease known as idiopathic pulmonary fibrosis. The study, published in the journal *Respirology*, found that patients with the condition who also cough are more likely to develop advanced forms of the disease that may be life threatening.

When idiopathic [pulmonary fibrosis](#) develops, tissue deep in the lungs becomes thick and scarred, likely due to a response to an unknown substance. The condition affects approximately 100,000 individuals in the [United States](#), and up to half die within three years of being diagnosed.

Almost all patients with idiopathic pulmonary fibrosis experience shortness of breath. The second most common symptom is cough. [Shortness of breath](#) is a known warning sign that a patient has a serious form of the disease, but little is known about the importance of cough. To investigate, Christopher Ryerson, MD, of the University of California, San Francisco, and his colleagues studied 242 patients with idiopathic pulmonary fibrosis. They found that cough was present in 84% of patients. It was more common in patients with advanced disease and in those who had never smoked. Also, the presence of cough predicted more rapid [disease progression](#), regardless of the severity of a patient's disease. The study's findings indicate that the presence of cough may predict which patients are likely to die prematurely or need a [lung](#) transplant in the near future.

To investigate, Christopher Ryerson, MD, of the University of California, San Francisco, and his colleagues studied 242 patients with idiopathic pulmonary fibrosis. They found that cough was present in 84% of patients. It was more common in patients with advanced disease and in those who had never smoked. Also, the presence of cough predicted more rapid disease progression, regardless of the severity of a patient's disease. The study's findings indicate that the presence of cough may predict which patients are likely to die prematurely or need a [lung transplant](#) in the near future.

The authors concluded that patients with idiopathic pulmonary fibrosis who cough may have a worse prognosis compared with patients who do not cough. While additional studies are needed to validate the results, patients who cough may benefit from closer monitoring and more aggressive treatments.

"These findings improve our understanding of cough in idiopathic pulmonary fibrosis," said Dr. Ryerson. "The reason for the association between cough and never having smoked is unknown, but may provide insight into the pathogenesis of cough in idiopathic pulmonary fibrosis and thus prompt future research in this area," he added.

Provided by Wiley

Citation: Cough may warn of danger for patients with lung-scarring disease (2011, October 18) retrieved 5 May 2024 from

<https://medicalxpress.com/news/2011-10-danger-patients-lung-scarring-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--