

# High-dose vitamin D may not be better than low-dose vitamin D in treating MS

October 24 2011

---

Low vitamin D levels are associated with an increased risk of developing multiple sclerosis (MS), but the first randomized, controlled trial using high-dose vitamin D in MS did not find any added benefit over and above ongoing low-dose vitamin D supplementation, according to a study published in the October 25, 2011, issue of *Neurology*, the medical journal of the American Academy of Neurology.

"We did not find added benefit from high-dose vitamin D over and above ongoing low-dose vitamin D supplementation, but these results need to be confirmed with larger studies," said Mark S. Stein, MBBS, PhD, FRACP, of The Royal Melbourne Hospital and The Walter and Eliza Hall Institute of Medical Research in Parkville, Australia.

The six-month study involved 23 people with the relapsing-remitting form of MS. All of the participants received low-dose vitamin D (1,000 international units daily) to prevent any [vitamin D deficiency](#). Half of the participants also received high-dose [vitamin D2](#) to elevate their blood vitamin D to high levels (with a target serum 25-hydroxyvitamin D level of 130-175nM). The other half received a placebo high-dose.

[MRI scans](#) of the participants' brains were performed at the start of the study and again after four, five and six months. There was no significant difference between the two groups in the number of new abnormalities that had formed in the brain after six months and no significant difference in the change in the total volume of [brain abnormalities](#).

Four of the 11 people taking the high-dose vitamin D, or 37 percent, had a [relapse](#) where their MS symptoms worsened during the study, while none of the 12 people taking only low-dose vitamin D had any relapses.

Stein noted that the study involved people who had MS for an average of six years. "It's possible that studies of high-dose vitamin D at an earlier stage of MS may lead to different results," he said.

Provided by American Academy of Neurology

Citation: High-dose vitamin D may not be better than low-dose vitamin D in treating MS (2011, October 24) retrieved 3 May 2024 from <https://medicalxpress.com/news/2011-10-high-dose-vitamin-d-low-dose-ms.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--