

Next-generation brain stimulation may improve treatment of Parkinson's disease

October 19 2011

Parkinson's disease (PD) is a devastating and incurable disease that causes abnormal poverty of movement, involuntary tremor, and lack of coordination. A technique called deep brain stimulation (DBS) is sometimes used to improve motor symptoms in patients with advanced disease. Now, a study published by Cell Press in the October 20 issue of the journal *Neuron* describes a new and more effective DBS paradigm that makes real-time adjustments in response to disease dynamics and progression and may be better for managing symptoms of advanced PD.

DBS involves implantation of a medical device that functions as a "[brain pacemaker](#)." Essentially, this device sends [electrical impulses](#) to specific regions of the brain and alters brain activity in those regions in a controlled manner. Although the underlying principles are not entirely clear, DBS has provided significant therapeutic benefits for movement disorders like PD and for affective disorders like chronic pain and [major depression](#).

After implantation of the DBS device, stimulation parameters, such as frequency and intensity of stimulation, must be programmed and adjusted over several months by a highly trained clinician. The goal is to maximize clinical improvement and minimize stimulation-induced side effects. Adjustments typically occur every 3 to 12 months when the patient visits the clinic, with the parameters remaining the same between visits. Unfortunately, this results in stimulation that does not keep up with the dynamic nature of PD.

"In recent years, the role of PD driven aberrant discharge patterns of [neuronal activity](#) have emerged as pivotal in the pathophysiology of the disease, and there is an urgent need for an automatic and dynamic system that can continually adjust the stimulus in response to ongoing [pathological changes](#)," explains Dr. Boris Rosin and Professor Hagai Bergman from The Hebrew University of Jerusalem. The authors tested several new paradigms for real-time adaptive DBS in a primate model of PD, in which the delivered stimulus was triggered by the ongoing brain activity.

The researchers discovered that real-time adaptive DBS paradigms alleviated PD motor symptoms and reduced abnormal neural activity more efficiently than standard DBS. The results provided new insight into brain activity underlying PD pathology and suggested that clinical improvement was achieved by disruption of a particular pattern from the variety of abnormal activity seen in the Parkinsonian brain. "It is our hope that in the near future we will see a new era of DBS strategies, based on real-time adaptive paradigms targeted at different pathological brain activity," conclude Dr. Rosin and Prof. Bergman. "Such strategies have potential not only for the treatment of PD, but perhaps other neurological disorders with a clear pathological pattern of [brain activity](#)."

Provided by Cell Press

Citation: Next-generation brain stimulation may improve treatment of Parkinson's disease (2011, October 19) retrieved 5 May 2024 from <https://medicalxpress.com/news/2011-10-next-generation-brain-treatment-parkinson-disease.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--