

# Study reports predictors of poor hand hygiene in an emergency department

October 3 2011

---

Researchers studying hand hygiene of healthcare workers in the emergency department found certain care situations, including bed location and type of healthcare worker performing care, resulted in poorer hand hygiene practice. The study was reported in the November issue of *Infection Control and Hospital Epidemiology*, the journal of the Society for Healthcare Epidemiology of America.

"We found that receiving care in a hallway bed was the strongest predictor of your healthcare providers not washing their hands," said Dr. Arjun Venkatesh, an emergency medicine resident at Brigham and Women's hospital in Boston, and author of the study. Dr. Venkatesh believes this finding will focus attention on infectious risks that are created by the national trend of emergency department crowding.

The study, which is the largest to date to evaluate hand hygiene in an emergency department, confirmed many known contributors of poor [hand washing](#) practices. For example, the researchers observed providers wearing gloves during patient care instead of washing their hands, an inappropriate substitution for [infection control](#) purposes.

The study also found that workers who transport patients between hospital departments and rooms were less likely to wash their hands compared to other [healthcare workers](#). This may be because these workers receive less training in [hand hygiene](#) procedures than other workers.

The researchers collected data on over 5,800 patient encounters in the emergency department. Overall, appropriate hand washing practices were used 90 percent of the time.

The authors hope the study will lay the foundation for future research and quality improvements in understanding the role of the emergency department in healthcare associated infections.

"With nearly one in five US residents visiting an [emergency department](#) each year, and emergency departments serving as a frequent interface between the public and patients with [communicable diseases](#), we have to build systems that ensure the highest standards of hand washing and infection control to ensure the safest care for all patients," concluded Dr. Jeremiah Schuur, Director for Quality, Safety and Performance Improvement for the Department of Emergency Medicine, Brigham and Women's Hospital.

Provided by Society for Healthcare Epidemiology of America

Citation: Study reports predictors of poor hand hygiene in an emergency department (2011, October 3) retrieved 26 April 2024 from <https://medicalxpress.com/news/2011-10-predictors-poor-hygiene-emergency-department.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--