

Anti-clotting drugs do not increase bleeding risk in GI procedure, study finds

November 1 2011

Patients with recent use of aspirin, nonsteroidal anti-inflammatory drugs (NSAIDs), or anti-clotting drugs such as clopidogreal (Plavix) do not appear to have an increased risk of bleeding during or after removal of precancerous lesions in the digestive tract, according to results of a Mayo Clinic study. The findings, culled from a review of 1,382 procedures of patients treated at Mayo Clinic in Florida, are being presented at the annual meeting of the American College of Gastroenterology in Washington, D.C.

"Patients who have recently used these medications worry a lot about bleeding when lesions are removed from their esophagus, stomach, or colon. This study shows that such concerns may not be warranted," says the lead author Bashar Qumseya, M.D.

"In the past, we told patients to discontinue these agents, if possible, in order to prevent bleeding. But now we know that this may not be necessary," says Dr. Qumseya, a second year gastroenterology fellow at Mayo Clinic Florida.

Patients with recent use of aspirin, nonsteroidal anti-inflammatory drugs ([NSAIDs](#)), or anti-clotting drugs such as clopidogreal ([Plavix](#)) do not appear to have an increased risk of bleeding during or after removal of [precancerous lesions](#) in the digestive tract, according to results of a Mayo Clinic study. The findings, culled from a review of 1,382 procedures of patients treated at Mayo Clinic in Florida, are being presented at the annual meeting of the American College of

Gastroenterology in Washington, D.C.

To conduct the study, Dr. Qumseya and his team examined Mayo Clinic's database of patients who received an endoscopic mucosal resection (EMR) from 1999 to 2010. An EMR is a minimally [invasive procedure](#) that can remove lesions from throughout the [GI tract](#).

The researchers found that bleeding during an EMR occurred in 3.9 percent of patients, and that it was linked to procedures in the esophagus and stomach. Bleeding that occurred days to weeks after the procedure was found in 2.7 percent of patients, and was associated with lesions anywhere in the [digestive tract](#) that were larger than 5 centimeters. Use of anti-inflammatory and anti-clotting agents, however, did not increase risk of bleeding either before or after an EMR.

"It is very important that patients know the risk of all medical procedures, and these findings will help us do the best job we can of informing our patients about those risks," Dr. Qumseya says.

Provided by Mayo Clinic

Citation: Anti-clotting drugs do not increase bleeding risk in GI procedure, study finds (2011, November 1) retrieved 20 April 2024 from <https://medicalxpress.com/news/2011-11-anti-clotting-drugs-gi-procedure.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.