

Time to test assumptions about health effects that guide risk assessment: toxicologist

November 21 2011

Governments and the nuclear industry have failed to address serious data gaps and untested assumptions guiding exposure limits to Cesium (Cs)-137 released in the Chernobyl accident in 1986 and this year's incident at Fukushima, says University of Massachusetts Amherst toxicologist Edward Calabrese. It's time now to move toward adopting more evidence-based risk assessment for the future, he adds.

"It is also critical that the linear, no-threshold (LNT) model and the alternative models, such as the threshold and hormesis models, be objectively assessed so that society can be guided by scientific data and validated models rather than ideological perspectives that stealthily infected the <u>risk assessment</u> process for <u>ionizing radiation</u> and <u>carcinogenic chemicals</u>," he states.

Calabrese's commentary, "Improving the scientific foundations for estimating health risks from the <u>Fukushima</u> incident," is included in this week's special issue of <u>Proceedings of the National Academy of</u> <u>Sciences</u> devoted to documenting the estimated release magnitude and distribution of Cs-137 from the nuclear incident in Japan after the March earthquake and tsunami.

After his career-long study of hormesis persuaded Calabrese that low doses of some chemicals and radiation are benign to humans or even helpful, he says a "comprehensive reappraisal" of the LNT model for cancer risk assessment is urgently needed. He argues that the LNT model was incorporated into U.S. regulatory policy in the 1950s based on faulty



assumptions and even manipulation of the scientific literature. Its use has become codified in regulatory agencies despite its "questionable scientific foundations," he says.

In addition to over-reliance on the LNT, Calabrese contends, regulators also place too much weight on assumptions about the ingestion of contaminants in soil by children, in particular dioxin, which date from the 1980s at Times Beach, Mo. "Subsequent soil ingestion studies in children would prove this default exposure assumption represented a massive overestimation, being too high by 200-fold," he points out. Without follow-up studies to provide more accurate data, "costs of clean up at Times Beach alone would have been many billions of dollars more," he adds, illustrating that non-validated assumptions can markedly affect the risk assessment outcome.

Calabrese also criticizes expert advisory groups and government agencies for rendering exposure guidance "based on hypothetical risks of Cs-137" and using highly precise estimates that give "a false impression of considerable accuracy." In fact, he says, acceptable levels of Cs-137 exposure in Japan are more than three times higher than levels permitted in the Ukraine, while both are probably based on "little independent analysis."

Overall, precautionary urges that pressure regulators to rely on the most conservative option have a downside, in Calabrese's opinion. They lead to multiplicative protective factors that can add substantially to remediation costs "without validated assurances of accompanying benefit."

Lower is not always better when it comes to enhancing public health, he says. "It is time for the responsible governmental and industrial organizations to develop a practical plan to fill important data gaps."



Provided by University of Massachusetts at Amherst

Citation: Time to test assumptions about health effects that guide risk assessment: toxicologist (2011, November 21) retrieved 2 May 2024 from https://medicalxpress.com/news/2011-11-assumptions-health-effects-toxicologist.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.