

# Body mass index associated with short-term mortality rates following surgery

November 21 2011

---

Body Mass Index (BMI) appears to be associated with 30-day mortality risk following surgical procedures, and patients with a BMI of less than 23.1 appear to be at highest risk of death, according to a report published Online First by *Archives of Surgery*.

"Recent reports suggest that the [prevalence of obesity](#) among U.S. adults has increased more than 100 percent since 1990," the authors write as background information in the article. "This analysis examines the relationship between obesity (as measured by [body mass index](#) [BMI; calculated as weight in kilograms divided by height in meters squared]) and [surgical mortality](#)."

Using data from the American College of Surgeons National Surgical [Quality Improvement Program](#), Florence E. Turrentine, Ph.D., R.N., and colleagues with the University of Virginia, Charlottesville, examined the relationship between BMI and 30-day mortality among 189,533 patients who underwent general or vascular surgical procedures at one of 183 sites in either 2005 or 2006.

Of the 189,533 patients included in the analysis, 3,245 (1.7 percent) died within 30 days of surgery. The authors found that the percentage of deaths (2.8 percent) among patients with a BMI less than 23.1 was over twice that of the percentage of deaths (1 percent) among patients with a BMI of 35.3 or higher. Additionally, patients with a BMI of less than 23.1 had statistically significant increased risk of death, with 40 percent higher odds of death than patients with mid-range BMI (between 26.3

and 29.6).

When examining data by procedure category, the authors also found that patients who underwent exploratory laparotomy had the highest percentage of death (13.9 percent) compared with patients in all other categories of principal surgery, and patients who underwent breast lumpectomy had one of the lowest overall mortality percentages (0.1 percent). The authors also found a statistically significant interaction between BMI and procedure category, indicating that the association between BMI and mortality was statistically different for patients who underwent these procedures (including colostomy, wound debridement, musculoskeletal system procedures, upper gastrointestinal procedures, colorectal resection, hernia repair, among others) compared with patients who underwent laparoscopy.

"These results indicate that BMI is a significant predictor of mortality within 30 days of surgery, even after adjusting for the contribution to [mortality risk](#) made by type of surgery and for a specific patient's overall expected risk of death," the authors conclude.

**More information:** *Arch Surg*. Published online November 21, 2011. [doi:10.1001/archsurg.2011.310](https://doi.org/10.1001/archsurg.2011.310)

Provided by JAMA and Archives Journals

Citation: Body mass index associated with short-term mortality rates following surgery (2011, November 21) retrieved 5 May 2024 from <https://medicalxpress.com/news/2011-11-body-mass-index-short-term-mortality.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.