


# How the brain strings words into sentences

November 24 2011, By Daniel Stolte

---



Using magnetic resonance imaging of the brain, researchers can visualize the two main language processing regions, Broca's region (yellow) and Wernicke's region (purple). (Image: Stephen Wilson)

(Medical Xpress) -- Distinct neural pathways are important for different aspects of language processing, researchers have discovered, studying patients with language impairments caused by neurodegenerative diseases.

While it has long been recognized that certain areas in the brain's left hemisphere enable us to understand and produce language, scientists are still figuring out exactly how those areas divvy up the highly complex processes necessary to comprehend and produce language.

Advances in brain imaging made within the last 10 years have revealed that highly complex [cognitive tasks](#) such as [language processing](#) rely not

only on particular regions of the [cerebral cortex](#), but also on the white matter fiber pathways that connect them.

"With this new technology, scientists started to realize that in the [language network](#), there are a lot more connecting pathways than we originally thought," said Stephen Wilson, who recently joined the University of Arizona's department of speech, language and hearing sciences as an assistant professor. "They are likely to have different functions because the brain is not just a homogeneous conglomerate of cells, but there hasn't been a lot of evidence as to what kind of information is carried on the different pathways."


Working in collaboration with his colleagues at the UA, the department of neurology at the University of California, San Francisco and the Scientific Institute and University Hospital San Raffaele in Milan, Italy, Wilson discovered that not only are the connecting pathways important for language processing, but they specialize in different tasks.

Two [brain areas](#) called Broca's region and Wernicke's region serve as the main computing hubs underlying language processing, with dense bundles of [nerve fibers](#) linking the two, much like [fiber optic cables](#) connecting [computer servers](#). But while it was known that Broca's and Wernicke's region are connected by upper and a lower white matter pathways, most research had focused on the nerve cells clustered inside the two language-processing regions themselves.

Working with patients suffering from language impairments because of a variety of neurodegenerative diseases, Wilsons' team used brain imaging and language tests to disentangle the roles played by the two pathways. Their findings are [published](#) in a recent issue of the scientific journal *Neuron*.

"If you have damage to the lower pathway, you have damage to the

lexicon and semantics," Wilson said. "You forget the name of things, you forget the meaning of words. But surprisingly, you're extremely good at constructing sentences."



Assistant professor Stephen Wilson studies how the brain processes language by combining brain imaging with performance-based language tests. (Photo: D. Stolte/UANews)

"With damage to the upper pathway, the opposite is true; patients name things quite well, they know the words, they can understand them, they can remember them, but when it comes to figuring out the meaning of a complex sentence, they are going to fail."

The study marks the first time it has been shown that upper and lower tracts play distinct functional roles in language processing, the authors write. Only the upper pathway plays a critical role in syntactic processing.

Wilson collected the data while he was a postdoctoral fellow working with patients with neurodegenerative diseases of varying severity, recruited through the Memory and Aging Center at UCSF. The study included 15 men and 12 women around the age of 66.

Unlike many other studies investigating acquired language disorders, which are called aphasias and usually caused by damage to the brain, Wilson's team had a unique opportunity to study patients with very specific and variable degrees of brain damage.

"Most aphasias are caused by strokes, and most of the strokes that affect language regions probably would affect both pathways," Wilson said. "In contrast, the patients with progressive aphasias who we worked with had very rare and very specific [neurodegenerative diseases](#) that selectively target different brain regions, allowing us to tease apart the contributions of the two pathways."

To find out which of the two nerve fiber bundles does what in language processing, the team combined magnetic resonance brain imaging technology to visualize damaged areas and language assessment tasks testing the participants' ability to comprehend and produce sentences.

"We would give the study participants a brief scenario and ask them to complete it with what comes naturally," Wilson said. "For example, if I said to you, 'A man was walking along the railway tracks. He didn't hear the train coming. What happened to the man?' Usually, you would say, 'He was hit by the train,' or something along those lines."

"But a patient with damage to the upper pathway might say something like 'train, man, hit.' We found that the lower pathway has a completely different function, which is in the meaning of single words."

To test for comprehension of the meaning of a sentence, the researchers presented the patient with a sentence like, "The girl who is pushing the boy is green," and then ask which of the two pictures depicted that scenario accurately.

"One picture would show a green girl pushing a boy, and the other would

show a girl pushing a green boy," Wilson said. "The colors will be the same, the agents will be the same, and the action is the same. The only difference is, which actor does the color apply to?"

"Those who have only lower pathway damage do really well on this, which shows that damage to that pathway doesn't interfere with your ability to use the little function words or the functional endings on words to figure out the relationships between the words in a sentence."

Wilson said that most previous studies linking neurodegeneration of specific regions with cognitive deficits have focused on damage to gray matter, rather than the [white matter](#) that connects regions to one another.

"Our study shows that the deficits in the ability to process sentences are above and beyond anything that could be explained by gray matter loss alone," Wilson added. "It is the first study to show that damage to one major pathway more than then other major pathway is associated with a specific deficit in one aspect of language."

**More information:** [www.cell.com/neuron/abstract/S0896-6273\(2011\)000835-X](http://www.cell.com/neuron/abstract/S0896-6273(2011)000835-X)

Provided by University of Arizona

Citation: How the brain strings words into sentences (2011, November 24) retrieved 10 April 2024 from <https://medicalxpress.com/news/2011-11-brain-words-sentences.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---