

People with early Alzheimer's disease may be more likely to have lower BMI

November 21 2011

Studies have shown that people who are overweight in middle age are more likely to develop Alzheimer's disease decades later than people at normal weight, yet researchers have also found that people in the earliest stages of Alzheimer's disease are more likely to have a lower body mass index (BMI). A current study examines this relationship between Alzheimer's disease and BMI.

The study, published in the November 22, 2011, print issue of *Neurology*, the medical journal of the American Academy of Neurology, examined 506 people with advanced brain imaging techniques and analyses of cerebrospinal fluid to look for biomarkers for Alzheimer's disease, which can be present years before the first symptoms begin. The participants, who were part of the Alzheimer's Disease Neuroimaging Initiative, included people with no memory problems, people with mild cognitive impairment, or mild [memory problems](#), and people with Alzheimer's disease.

The study found that in people with no memory or thinking problems and in people with mild cognitive impairment, those who had the Alzheimer's biomarkers were also more likely to have a lower BMI than those who did not have the biomarkers.

For example, 85 percent of the people with mild cognitive impairment who had a BMI below 25 had signs of the beta-amyloid plaques in their brains that are a hallmark of the disease, compared to 48 percent of those with [mild cognitive impairment](#) who were overweight. The

relationship was also found in people with no memory or thinking problems.

"These results suggest Alzheimer's disease [brain changes](#) are associated with systemic [metabolic changes](#) in the very earliest phases of the disease," said study author Jeffrey M. Burns, MD, MS, of the University of Kansas School of Medicine in Kansas City and a member of the American Academy of Neurology. "This might be due to damage in the area of the brain called the hypothalamus that plays a role in regulating [energy metabolism](#) and food intake. Further studies should investigate whether this relationship reflects a systemic response to an unrecognized disease or a long-standing trait that predisposes a person to developing the disease."

Provided by American Academy of Neurology

Citation: People with early Alzheimer's disease may be more likely to have lower BMI (2011, November 21) retrieved 24 April 2024 from <https://medicalxpress.com/news/2011-11-people-early-alzheimer-disease-bmi.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--