

Psoriasis is associated with impaired HDL function

November 16 2011

Collaborative research from Perelman School of Medicine at the University of Pennsylvania has shown that psoriasis patients have an increased risk of heart attack, stroke and cardiovascular death, especially if the psoriasis is moderate to severe. Now, Penn researchers have discovered the potential underlying mechanism by which the inflammatory skin disease impacts cardiovascular health. In two new studies presented at the 2011 American Heart Association Scientific Sessions, Penn researchers show that the systemic inflammatory impact of psoriasis may alter both the makeup of cholesterol particles and numbers, as well as impair the function of high density lipoprotein (HDL), the "good" cholesterol.

"Anecdotally, many researchers have observed that HDL levels may be lower in states of inflammation, such as [rheumatoid arthritis](#), [psoriasis](#) and even obesity," said lead study author Nehal Mehta, MD, MSCE, director of Inflammatory Risk in Preventive Cardiology at Penn.

"However, these new findings suggest that in addition to lower levels, [chronic inflammation](#) associated with conditions like psoriasis may change the composition and decrease the function of HDL as well."

In the current studies, researchers enrolled 78 patients with psoriasis and 84 control subjects. In the first study, the authors measured fasting [lipid levels](#) and examined the number and size of cholesterol particles using [nuclear magnetic resonance](#) (NMR) spectroscopy. This analysis revealed that patients with psoriasis had a higher number of smaller LDL particles, or "bad" cholesterol, which was independent of traditional risk

factors and obesity. "It was striking that the NMR profiles from patients with psoriasis resembled those seen in patients with diabetes, and that these patients with psoriasis had otherwise normal traditional lipid panels" Dr. Mehta added.

In the second study, the researchers measured HDL efflux, which is the ability of a patient's HDL to remove cholesterol from cells involved in atherosclerosis. This process, known as 'reverse cholesterol transport', is why HDL may have protective properties. In a previous study, researchers at Penn have demonstrated that measuring HDL efflux capacity may be a more effective barometer of protection from heart disease than measuring HDL levels alone.

In this same group of patients who had normal cholesterol levels compared to controls, patients with psoriasis demonstrated dramatically reduced HDL efflux capacity compared to control patients. This negative association observed between psoriasis and HDL efflux persisted after adjusting for traditional lipid levels and other traditional risk factors, including body mass index (BMI).

"Patients with psoriasis had an approximate 25 percent reduction in the HDL efflux capacity than the controls, despite their relatively normal overall lipid profiles which leads to the question of whether function is more important than concentration in chronic inflammatory states" Dr. Mehta noted.

The new findings may provide a critical clue to the link between psoriasis and heart disease, but the researchers say larger studies are needed to validate their findings. Joel M. Gelfand, MD, MSCE, assistant professor of Dermatology and Epidemiology, and a senior author on the studies, said "We've been able to show that psoriasis is an important risk factor for vascular disease, and now we may finally be able to identify and ultimately treat the pathways by which psoriasis increases these

risks."

Provided by University of Pennsylvania School of Medicine

Citation: Psoriasis is associated with impaired HDL function (2011, November 16) retrieved 11 May 2024 from <https://medicalxpress.com/news/2011-11-psoriasis-impaired-hdl-function.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.