

Researchers uncover steps in synapse building, pruning

November 16 2011

Like a gardener who stakes some plants and weeds out others, the brain is constantly building networks of synapses, while pruning out redundant or unneeded synapses. Researchers at The Jackson Laboratory led by Assistant Professor Zhong-wei Zhang, Ph.D., have discovered a factor in synapse-building, also showing that the building and pruning processes occur independent of each other.

Mammals are born with functioning but not-yet-developed brains. After birth, external stimuli and internal programs continue to shape the connections between neurons, known as [synapses](#), and the formation of networks of synapses known as [neuronal circuits](#). Some grow stronger, some grow weaker, redundant connections are eliminated, and so on. Such "plasticity," the ongoing refinement of [neural connections](#) and networks, continues throughout life, albeit more subtly with time and maturation.

Much about plasticity remains unknown. How the [neural circuits](#) are modified, what controls the modification, the mechanics of strengthening or eliminating specific synapses and much more are subjects of ongoing research. Besides gaining a better picture of normal brain development, scientists seek to understand the errors in synapse building and pruning that are associated with autism, mental retardation and schizophrenia.

Zhang and colleagues investigated a major type of synapse in the brain (called the glutamatergic synapse) that undergoes rapid refinement soon

after birth. What they discovered is that these synapses are strengthened through the addition of a particular kind of glutamate receptors, beginning about a week after birth for mice. Notably, sensory deprivation disrupts the strengthening of the synapses, highlighting the role of early experience in synapse building.

In a somewhat surprising finding, the Zhang lab also discovered that the elimination of redundant synapses was not dependent on the other synapses' being strengthened. Since synaptic strengthening usually precedes removal of redundant synapses, it was not known if such elimination is dependent on the prior strengthening. In mice lacking the receptor, which prevented significant strengthening of synaptic connections, redundant synapses were eliminated as usual.

More information: Wang et al.: Elimination of redundant synaptic inputs in the absence of synaptic strengthening. *Journal of Neuroscience*, Nov. 16, 2011, [DOI:10.1523/JNEUROSCI.4569-11.2011](https://doi.org/10.1523/JNEUROSCI.4569-11.2011)

Provided by Jackson Laboratory

Citation: Researchers uncover steps in synapse building, pruning (2011, November 16) retrieved 26 April 2024 from <https://medicalxpress.com/news/2011-11-uncover-synapse-pruning.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|