

Association of cognitive function in adolescence and subsequent risk of subdural hematoma

December 27 2011

Anna Nordström and Peter Nordström of Umea University, Sweden, report their analysis of a prospective nationwide cohort of 440,742 Swedish men in this week's *PLoS Medicine*, finding that reduced cognitive function in young adulthood was associated with increased risk of subdural hematoma later in life, whereas a higher level of education and physical fitness were associated with a decreased risk.

These results need to be confirmed in other large studies.

The authors say: "An exploration of the mechanistic basis for these associations might allow the construction of public health interventions aimed at reducing the population incidence of [subdural hematoma]... The present study was observational and thus inferences about causality should [...] be made with great caution. However, some of the findings in the present study may suggest a cause–effect relationship."

More information: Nordström A, Nordström P (2011) Cognitive Performance in Late Adolescence and the Subsequent Risk of Subdural Hematoma: An Observational Study of a Prospective Nationwide Cohort. *PLoS Med* 8(12): e1001151. [doi:10.1371/journal.pmed.1001151](https://doi.org/10.1371/journal.pmed.1001151)

Provided by Public Library of Science

Citation: Association of cognitive function in adolescence and subsequent risk of subdural hematoma (2011, December 27) retrieved 23 April 2024 from <https://medicalxpress.com/news/2011-12-association-cognitive-function-adolescence-subsequent.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.