

Common bone drug may extend life of replacement joints

December 7 2011

People who take bisphosphonates after joint replacement surgery are less likely to need a repeat operation, finds a new study published in the British Medical Journal today.

While hip and knee replacements are cost effective procedures, substantial numbers of patients require revision [surgery](#) within 10 years to replace the implant because of infection, wear, loosening, or other mechanical failures.

Oral bisphosphonates are used to prevent fractures and to treat common bone diseases like osteoporosis. Some experts think that bisphosphonates may improve the survival time of replacement joints through their effects on reducing [bone resorption](#), but evidence is scarce and controversial.

So a team of researchers, led by Professor Nigel Arden at the University of Oxford, set out to test whether bisphosphonate use is related to improved implant survival time.

Using data from the General Practice Research Database, which contains anonymised data on three million UK primary care patients, they identified all patients over the age of 40 undergoing hip or [knee replacement surgery](#) from 1986-2006 for osteoarthritis.

Patients who were prescribed bisphosphonates at least six months before revision surgery were defined as bisphosphonate users. Non-users were

defined as those who were never prescribed bisphosphonates or who had their first prescription after revision surgery.

Factors that could affect the results, such as age, sex, [body mass index](#) (BMI), smoking status, [alcohol intake](#), use of calcium and vitamin D supplements, and previous fractures were also taken into account.

Patients were tracked for a maximum of 15 years after their operation to calculate the rate of revision surgery.

Bisphosphonate users had a lower rate of revision rate at five years than non-users. Bisphosphonate use was associated with an almost twofold increase in implant survival time.

The researchers estimate that, for every 107 joint replacement patients taking bisphosphonates, one revision operation would be avoided.

And they suggest that the results could be due to bisphosphonate therapy suppressing the long term inflammatory response around the implant after surgery that often leads to bone loss and loosening of the implant.

Further studies are needed to confirm these findings, write the authors. However, they say that the protective effect of [bisphosphonates](#) seen in this study can be assumed to be valid for the wider population of patients undergoing total hip or knee replacement in the UK.

Provided by British Medical Journal

Citation: Common bone drug may extend life of replacement joints (2011, December 7) retrieved 11 May 2024 from <https://medicalxpress.com/news/2011-12-common-bone-drug-life-joints.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.