

Study suggests feelings of guilt may be a top factor in PTSD

December 6 2011, By Gregg Zoroya

A leading cause of post-traumatic stress disorder is guilt that troops experience because of moral dilemmas faced in combat, according to preliminary findings of a study of active-duty Marines.

The conflicts that servicemembers feel may include "survivor's guilt," from living through an attack in which other servicemembers died, and witnessing or participating in the unintentional killing of women or children, researchers involved in the study say.

"How do they come to terms with that? They have to forgive themselves for pulling the trigger," says retired Navy captain Bill Nash, a psychiatrist and study co-author.

The idea of "moral injury" as a cause of [PTSD](#) is new to psychiatry. The [American Psychiatric Association](#) is only now considering new [diagnostic criteria](#) for the disorder that would include feelings of shame and guilt, says David Spiegel, a member of the working group rewriting the PTSD section.

Traditionally, PTSD symptoms such as nightmares or numbness to the world have been linked to combat violence, fear of being killed or loss of friends.

Half of all Iraq and Afghanistan veterans treated by the Department of Veterans Affairs have been diagnosed with [mental health issues](#) and the most common is PTSD, which is experienced by nearly 200,000 of these

veterans, according to the VA.

PTSD caused by moral injury can lead to more severe reactions such as [family violence](#) or even suicide, says Jonathan Shay, a psychiatrist who has worked on military mental health policies.

The Marine Corps study helps expand the knowledge of the relationship between moral injury and PTSD, says Shira Maguen, a psychologist and VA researcher who has studied links between killing and the disorder among Vietnam War, Gulf War and Iraq War veterans.

"This (Marine Corps) study is important because so little work has been done to understand moral injury in a scientific context," Maguen says.

The ongoing research involves about 2,600 Marines and sailors examined before and after combat tours.

The preliminary findings on moral injury were gleaned from 208 Marines involved in severe combat in Afghanistan in 2009 and 2010. It showed that three months after coming home, 7 percent of the Marines likely had PTSD. Their condition was more closely linked to an inner conflict rather than threats to their lives, the sight of bodies or blood or family problems, the study said.

(c)2011 USA Today
Distributed by MCT Information Services

Citation: Study suggests feelings of guilt may be a top factor in PTSD (2011, December 6)
retrieved 18 April 2024 from <https://medicalxpress.com/news/2011-12-guilt-factor-ptsd.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--