

'Mindfulness' exercises help curb stress and fatigue associated with arthritis

December 20 2011

"Mindfulness" exercises, which focus on experiencing the present moment, no matter how difficult, can help curb the stress and fatigue associated with painful rheumatoid joint disease, indicates a small study published online in the *Annals of Rheumatic Diseases*.

The authors base their findings on 73 patients between the ages of 20 and 70, all of whom had had painful joint disease, caused by [rheumatoid arthritis](#), ankylosing [spondylitis](#), or psoriatic [arthritis](#) for at least a year.

Half of them were randomly allocated to scheduled "mindfulness" exercises, which took place in 10 group sessions over a period of 15 weeks, plus a booster session around six months after the course had completed.

The sessions, which were facilitated by [healthcare professionals](#) trained in mindfulness techniques, addressed particular topics, such as recognising individual limitations, and strong emotions, such as anger, joy, and sorrow.

The exercises, which were part of the Vitality [Training Programme](#) of VTP, encouraged participants to become aware of, and deliberately concentrate on their feelings, thoughts and bodily experiences, including pain, without judging or trying to avoid them.

Participants were also given creative exercises, such as guided imagery, music and drawing, and shared their experiences with other members of

the group.

The rest of the volunteers randomly allocated to the comparison group were given standard care plus a CD containing similar exercises for use at home, as and when they wanted.

[Stress levels](#), coping abilities, and symptom control, including pain and [fatigue](#), were assessed, using validated scores, immediately after all 10 sessions had finished, and again 12 months later.

In total, 67 participants completed all the assessments. These showed no differences in pain levels, disease activity or the ability to talk about feelings.

But there were significant differences in [levels of stress](#) and fatigue.

The number of participants with a high stress score of above 23 in the GHQ-20 questionnaire fell from 13 at the start of the study to two, just 12 months after the sessions had finished. Comparable figures in the comparison group were 10 and eight, respectively.

There was, however, a tangible fall in measured levels of fatigue in the intervention group: no such change was evident in the comparison group.

There have been previous attempts to use psychological and educational tactics to help people with arthritis cope better with the distressing aspects of the disease, but they have tended to be short term, say the authors.

The lasting improvements found with the VTP course "indicate that the participants may have incorporated some mindfulness strategies into their daily lives and that these strategies have strengthened their ability to respond to their stressful experience in a more flexible way," they say.

The authors emphasise that while the treatment of rheumatoid arthritis has improved greatly, it is less effective in those with more established disease, and that ultimately the disease can only be partly controlled, forcing many patients to make very demanding lifestyle changes.

"There is therefore a need for complementary interventions that enhance individuals' health-promoting resources and help them adjust to their disease," they conclude.

More information: To be published at ard.bmj.com

Provided by British Medical Journal

Citation: 'Mindfulness' exercises help curb stress and fatigue associated with arthritis (2011, December 20) retrieved 24 April 2024 from <https://medicalxpress.com/news/2011-12-mindfulness-curb-stress-fatigue-arthritis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.