

# Ovarian cancer study proves drug delays disease progression, may improve survival

December 28 2011

---

Treating ovarian cancer with the drug bevacizumab ("Avastin") delays the disease and may also improve survival, show the results of an international clinical trial co-led by Drs. Amit Oza of the Princess Margaret Cancer Program, University Health Network and Timothy Perren, St James's Institute of Oncology, Leeds, UK.

The findings, published today in the [New England Journal of Medicine](#), report that the drug halted the cancer's return for two months overall. However, for women with the highest risk disease, the delay was five to six months and in this group, the findings also indicate a strong trend to improved overall survival, which is being analysed until 2013.

"This is the first new drug in ovarian cancer in 15 years to improve outcome and I believe it should be considered as a potential new standard of care," says Dr. Oza, a medical oncologist who leads the Cancer Clinical Research Unit at Princess Margaret Hospital. He is also co-director of the hospital's Bras Family Drug Development Program and Professor, Faculty of Medicine, University of Toronto.

The seven-year study began in 2004 and enrolled 1,528 women with ovarian cancer at 263 centres, including 20 in Canada. [Avastin](#) was added to chemotherapy treatment and given intravenously every three weeks for 12 months.

The [drug](#) blocks growth factors that promote new blood vessels formation in tumours, thereby "starving" the cancer. It is not a cure,

explains Dr. Oza, but has a proven track record in delaying disease progression in other [types of cancer](#) including colorectal, lung, breast, kidney and brain.

"We now know that using Avastin in [ovarian cancer](#) for even this short time improves outcomes," says Dr. Oza. "The next step is to determine if giving it for a longer period would be of even greater benefit."

Similar findings from a U.S. study are also reported in this issue of the journal. Dr. Oza says the major difference between the two studies is that the women in the American study were given twice as much Avastin. "So the question now is would half the dose for double the duration improve outcomes even more? This is an area to investigate further."

Provided by University Health Network

Citation: Ovarian cancer study proves drug delays disease progression, may improve survival (2011, December 28) retrieved 27 April 2024 from <https://medicalxpress.com/news/2011-12-ovarian-cancer-drug-disease-survival.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--