

UGA study documents lung function declines in firefighters working at prescribed burns

December 7 2011



After monitoring firefighters working at prescribed burns in the southeastern U.S., University of Georgia researchers found that lung function decreased with successive days of exposure to smoke and other particulate matter.

"What we found suggested a decline in lung function across work seasons," said Olorunfemi Adetona, a postdoctoral research associate and lead author of the study published recently in the journal *Inhalation Toxicology*.

Luke Naeher, senior author and associate professor in the UGA College of Public Health, explained that the study was designed to investigate

whether the 26 [firefighters](#) experienced a decrease in lung function working at [prescribed burns](#) compared with days they spent away from the fires. Previously, researchers had looked only at changes in lung function of wildland firefighters on days with exposure to smoke.

"Over a 10-week season, these workers' respiratory functions slowly declined," Naeher said, adding that there is need to investigate the degree to which these declines returned to their baseline after the burn season. Although results of the study show that lung function at the start of two burn seasons in a limited number of nine firefighters in 2003 and 2004 did not vary significantly, more definitive answers relating to the issue of longer term effect of exposure on [lung function](#) would require a different study design.

In recent years, the U.S. [Forest Service](#) has sought to better understand and improve its occupational exposure limits for firefighters across the country. Most studies have concentrated on burns in Western states where exposure to and composition of wood-smoke [particulate matter](#) may vary to some degree when compared with fires in the Southeast, including South Carolina, where the study was done.

Naeher said the study provides some preliminary information regarding the [health effects](#) of [fine particulate matter](#) exposure that is intermediate between two exposure extremes. On the low extreme lie ambient air levels typical for developed countries, while inhalation of particles by a smoker represents the opposite extreme. Much research in the field has focused on health effects at both extremes. However, the study of exposure at intermediate levels, like that experienced by wildland firefighters, and women and children exposed to indoor air pollution from cook stoves in developing countries is limited. Naeher's research focuses on these two different populations, and he explains that the study of the body's response to these intermediate exposures may now be more urgent. For example, Naeher said, an initiative led by the United Nations

Foundation aims to put clean-burning cooking stoves in 100 million homes in developing countries by 2020.

Provided by University of Georgia

Citation: UGA study documents lung function declines in firefighters working at prescribed burns (2011, December 7) retrieved 6 May 2024 from

<https://medicalxpress.com/news/2011-12-uga-documents-lung-function-declines.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.