

Researchers suggest unconventional approach to control HIV epidemics

December 7 2011, By Mark Wheeler

(Medical Xpress) -- A new weapon has emerged to prevent HIV infection. Called pre-exposure prophylaxis, or PrEP, it is a strategy of providing medications to at-risk people before they are exposed to the virus. Having shown great promise in recent phase 3 clinical trials, PrEP may soon be rolled out for public use.

Because PrEP is based on the same drugs used to treat HIV-infected individuals, the big public health fear is that the dual use of these drugs will lead to skyrocketing levels of [drug resistance](#). But in a new study, UCLA researchers say the exact opposite is likely to happen.

Sally Blower, director of the UCLA Center for Biomedical Modeling and a professor at the Semel Institute for Neuroscience and Human Behavior at UCLA, and colleagues used sophisticated computer modeling to determine that a PrEP [prevention program](#) used alone, or current [HIV treatment](#) programs used alone, could indeed, separately, increase drug resistance. But if used together, the researchers say, resistance is likely to decrease.

Their findings appear in the current online edition of the peer-reviewed journal *Scientific Reports*, published by Nature Publishing Group.

"This was a very big surprise," said Blower, the study's senior author.

"We found that this counterintuitive effect will only occur if adherence to the PrEP prevention program, where individuals have to take a daily pill, is very high. This counterintuitive effect occurs when the beneficial

effect of PrEP in preventing infections is so great that it overcomes both its own detrimental effect on increasing resistance and the detrimental effect of current HIV treatments on increasing resistance."

Africa is ground zero for HIV and AIDS, a continent where the death rate is simply "awful," Blower said. Since the country of Botswana was one of the sites for the PrEP drug trial, Blower and her colleagues chose it for their modeling. Botswana has the best [health care system](#) in Africa, they said, yet 30 percent of women and 20 percent of men are infected with HIV.

"Botswana is likely to lead the way in rolling out PrEP," said Virginie Supervie, first author of the current study and a former graduate student in Blower's laboratory. "So officials there are worried about increasing levels of resistance."

Most health officials feel that if you have a good treatment program in place, it makes sense to establish a good prevention program in the same place. Health officials plan to implement PrEP prevention programs only where [treatment programs](#) are highly successful and levels of resistance are low, the researchers said.

But in a second counterintuitive finding, the UCLA researchers say this conventional approach is actually the worst strategy. Instead, they suggest, PrEP programs should be rolled out around treatment programs that are having little success and where rates of resistance are high. Their model shows that this unconventional approach would prevent the maximum number of infections and result in the greatest decrease in drug resistance.

"By cutting down infections, the PrEP programs will decrease the number entering treatment programs, and therefore, fewer individuals will acquire drug resistance," Blower said. "So introducing PrEP around

the worst treatment programs will have the most impact on reducing resistance."

A PrEP clinical trial that involved men who had sex with men and transgender women who had sex with men found that PrEP reduced the risk of acquiring [HIV infection](#) by 44 percent. Two other PrEP trials, which involved heterosexual men and women, showed significant reductions in risk, ranging as high as 73 percent.

"These results are very promising," Supervie said. "Our model shows that if the roll-out of PrEP is carefully planned, it could decrease resistance and increase the sustainability of treatment programs. But if it is not, resistance could increase, and the sustainability of treatment programs in resource-constrained countries could be compromised."

"The model we have designed is a very important health policy and planning tool," Blower said. "It can be used for any African country to decide where to roll out PrEP, as well as to predict the impact of PrEP on reducing their HIV epidemic. It is a critical time for decision-makers in Botswana and other African countries."

Other authors on the study included Meagan Barrett from UCLA; James S. Kahn from the University of California, San Francisco; and Godfrey Musuka, Themba L Moeti and Lesogo Busang from the African Comprehensive [HIV/AIDS Partnerships](#) in Gaborone, Botswana.

Provided by University of California Los Angeles

Citation: Researchers suggest unconventional approach to control HIV epidemics (2011, December 7) retrieved 1 May 2024 from <https://medicalxpress.com/news/2011-12-unconventional-approach-hiv-epidemics.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.