

In the brain, an earlier sign of autism

January 26 2012

In their first year of life, babies who will go on to develop autism already show different brain responses when someone looks at or away from them. Although the researchers are careful to say that the study, reported online on January 26 in the Cell Press journal *Current Biology*, is only a first step toward earlier diagnosis, the findings do suggest that direct brain measures might help to predict the future development of autism symptoms in infants as young as six months.

"Our findings demonstrate for the first time that direct measures of brain functioning during the first year of life associate with a later diagnosis of [autism](#)—well before the emergence of behavioral symptoms," said Mark Johnson of Birkbeck College, University of London.

The behaviors characteristic of autism emerge over the first few years of life, and firm diagnoses are now made in children only after the age of two. As a result, the vast majority of research on autism has necessarily concentrated on children two and older, who have already been diagnosed.

"We still know very little about the earliest appearing symptoms and warning signs," Johnson said.

To find out more, his team looked to six- to ten-month-old [babies](#) at greater risk of developing autism later in development because they had an older brother or sister with the condition. The researchers used passive sensors placed on the scalp to register brain activity while the

babies viewed faces that switched from looking at them to looking away from them or vice versa.

Earlier studies have shown that the human brain shows characteristic patterns of activity in response to eye contact with another person. That response is a critical foundation for face-to-face social interactions, and it is well known that older children diagnosed with autism show unusual patterns of eye contact and of [brain responses](#) to social interactions that involve eye contact.

The new studies reveal that the brains of infants who will go on to develop autism already process social information in a different way. "At this age, no behavioral markers of autism are yet evident, and so measurements of brain function may be a more sensitive indicator of risk," Johnson said.

It is important to note, however, that there were cases in which individual babies who showed these differences in [brain](#) function were not later diagnosed and vice versa. In other words, the method would require further refinement, most likely in combination with other factors, to form the basis of a predictor accurate enough for clinical use.

"Differences in the use of eye gaze to regulate social interaction is already a well-recognized early feature in many children with autism from the second year of life, and at present, it is these increasingly well-documented 'first signs' that will alert parents and professionals to possible differences," said Tony Charman from the Institute of Education, London, who co-led the study. "Future studies will be required to determine whether measurements of [brain function](#) such as those used in our study might one day play a role in helping to identify children at an even earlier age."

It will also be important to explore factors that might "protect" some

infants who do show early differences in their brains' responses to [eye contact](#) from going on to a diagnosis of autism, he added.

Provided by Cell Press

Citation: In the brain, an earlier sign of autism (2012, January 26) retrieved 2 May 2024 from <https://medicalxpress.com/news/2012-01-brain-earlier-autism.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.