

Starting smoking cessation medication earlier may make it easier to quit

January 5 2012

Smokers planning to kick the habit may have more success if they begin using a cessation medication several weeks before they actually try to quit. Those are the results of a clinical trial conducted by researchers at the University at Buffalo Roswell Park Cancer Institute (RPCI) and other institutions published recently in *Clinical Pharmacology and Therapeutics*.

The study focused on 35 women and 25 men, all smokers from Western New York who were on average 48 years old and smoked a pack of cigarettes per day. Participants who were randomized to take the smoking cessation medication varenicline (marketed as Chantix) for four weeks prior to trying to [quit smoking](#) were more likely to successfully quit smoking than those who took varenicline for just one week before quitting, which is the current standard therapy for the drug. Everyone took the medication for an additional 11 weeks after the quit day.

"Varenicline was designed to make smoking less rewarding, and our data suggests that it does that better when people take it for a few extra weeks before quitting," says Larry W. Hawk Jr., PhD, lead author and associate professor of psychology in the UB College of Arts and Sciences "If this finding holds up in larger studies, it could have a major impact on public health."

"We saw nearly full compliance, which suggests that this is not only a well-tolerated therapy, but one people can realistically stick with," says co-author Martin C. Mahoney, MD, PhD, associate professor of

oncology in RPCI's Departments of Medicine and [Health Behavior](#) and clinical associate professor in UB's School of Public Health and Health Professions and School of Medicine and Biomedical Sciences.

While many participants reported mild nausea, Mahoney says, the researchers found that the symptoms typically dissipated after a couple of weeks and may have helped reduce their desire to smoke.

"Whether through changes in taste or nausea, it seems this extra varenicline reduces smoking rates before people try to quit," Hawk says. "These changes should make it easier to quit smoking, but we also know that it takes some period of time for this new learning to occur. That's why we decided to see if a longer period of treatment with varenicline *before* smokers tried to quit would result in better outcomes, and it did in this small study."

Of special interest was the fact that women who took varenicline for four weeks were especially likely to reduce their smoking, possibly because they reported more nausea in the pre-quit period. After three weeks of treatments with varenicline, women reduced their smoking by more than 50 percent, on average. The men who took the varenicline for four weeks reduced their smoking by 26 percent. The researchers say that much larger studies are needed to tell whether the gender differences are real.

"This study suggests we may be able to take the most effective [smoking-cessation](#) treatment we have and make it work 50 percent better, just by giving the medication for a few weeks before smokers attempt to quit," concludes Hawk.

Provided by University at Buffalo

Citation: Starting smoking cessation medication earlier may make it easier to quit (2012, January 5) retrieved 19 April 2024 from

<https://medicalxpress.com/news/2012-01-cessation-medication-earlier-easier.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.