

Report: Mental illness struck 1 in 5 US adults in 2010

January 20 2012, By Jeannine Stein

One in five adults in the U.S. had a mental illness in 2010, with people ages 18 to 25 having the highest rates, according to a national survey.

The report from the Mental Health Services Administration's National Survey on <u>Drug Use</u> and Health, released Thursday, includes information from 68,487 completed surveys about <u>mental illness</u> (as defined by the American Psychiatric Association's Diagnostic and Statistical Manual of Mental Disorders-IV) and <u>substance abuse</u> among adults and children. Rates have remained fairly stable since 2009, with only a slight uptick in overall numbers.

Among the highlights, people in the 50-plus age bracket had the lowest incidence of any mental illness (14.3 percent), while those ages 18 to 25 had the highest, at 29.9 percent. Women had higher rates than men: 23 percent versus 16.8 percent

When broken down by racial and ethnic groups, the highest rates of mental illness were seen among people who reported two or more races (25.4 percent), followed by whites (20.6 percent), blacks (19.7 percent), Native Americans or Alaska natives (18.7 percent), Hispanics (18.3 percent) and <u>Asians</u> (15.8 percent).

About 5 percent of adults in the U.S. had a serious mental illness in 2010, a slightly higher number compared with 2009 (4.8 percent). That 18- to 25-year-old age group again had the highest incidence at 7.7 percent, and those 50 and older had the lowest at 3.2 percent. Women



again outpaced men, 6.5 percent to 3.4 percent. Those with a serious mental illness have trouble functioning, which affects their key life activities.

When broken down by racial and ethnic groups, the highest rates of serious mental illness occurred among adults who reported two or more races (9.3 percent), followed by Native Americans or Alaska natives (8.5 percent), whites (5.2 percent), <u>Hispanics</u> (4.6 percent), blacks (4.4 percent), Asians (2.6 percent) and <u>native Hawaiians</u> or other <u>Pacific Islanders</u> (1.6 percent).

"Mental illness is not an isolated public health problem," Substance Abuse and Mental Health Services administrator Pamela Hyde said in a news release. "Cardiovascular disease, diabetes, and obesity often coexist with mental illness and treatment of the mental illness can reduce the effects of these disorders."

The report also found that 39 percent of people who had any mental illness obtained <u>mental health services</u>, and among those with a serious mental illness, 60.8 percent received help.

Mental illness can lead to thoughts of suicide, and in 2010 8.7 million people thought seriously about taking their own lives. In that group, 2.5 million people made plans to kill themselves, and 1.1 million attempted suicide.

Substance abuse sometimes goes hand in hand with mental illness; those who had any mental illness or a serious mental illness had much higher substance abuse or dependence rates (20 percent) compared with those who didn't have a mental illness (6.1 percent).

Depression was an issue among children ages 12 to 17. Almost 2 million people in that age bracket had a major depressive episode in 2010 that



lasted at least two weeks. Those who had such an episode had more than twice the rate of using illicit drugs than their peers who didn't have a period of severe depression.

"Today's report ... provides further evidence that we need to continue efforts to monitor levels of mental illness in the United States in order to effectively prevent this important public health problem and its negative impact on total health," Ileana Arias of the Centers for Disease Control and Prevention said in the release.

(c)2012 the Los Angeles Times Distributed by MCT Information Services

Citation: Report: Mental illness struck 1 in 5 US adults in 2010 (2012, January 20) retrieved 3 May 2024 from <u>https://medicalxpress.com/news/2012-01-mental-illness-struck-adults.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.