


Researchers quantify muscle soreness

January 23 2012



This is a typical infrared image of a subject's arm before exercising. Credit: (c) Journal of Visualized Experiments

Quantifying how sore a person is after a long workout is a challenge for doctors and researchers, but scientists from Loma Linda and Asuza Pacific Universities think they may have figured it out. Their research article describing a new technique to measure muscle soreness will be published in the *Journal of Visualized Experiments (JoVE)*.

Delayed onset [muscle soreness](#) (DOMS) or exercise-induced [muscle damage](#) (EIMD) is one of the most common sports injuries, but without a reliable method of quantifying muscle soreness, assessing treatments is difficult.

Traditionally, muscle soreness has been measured using the visual

analogue scale (VAS). Participants mark their level of agreement to a statement along a continuous line. Rather than measuring soreness subjectively, the researchers used [thermal imaging](#) to detect subtle changes in the temperature of the skin above exercised muscles.

"The main advantage of this technique," said paper author Dr. Jerrold Petrofsky, "is that unlike visual scales, which are kind of a subjective measure of whether someone is sore or not, this technique gives you quantifiable, absolute data."

"There is no gold standard for measuring DOMS and other techniques, such as [needle biopsies](#), are invasive and painful for patients," said *JoVE* Editor, Leiam Colbert. "The technique presented here allows for earlier diagnosis and quicker treatment of soreness."

Provided by The Journal of Visualized Experiments

Citation: Researchers quantify muscle soreness (2012, January 23) retrieved 30 April 2024 from <https://medicalxpress.com/news/2012-01-quantify-muscle-soreness.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
