

Antidepressant use linked with less patient satisfaction after hip replacement

February 8 2012

Patients taking antidepressants up to three years prior to undergoing a total hip replacement (THR) were more likely to report greater pain before and after surgery and less satisfaction with their procedure, according to new research presented today at the 2012 Annual Meeting of the American Academy of Orthopaedic Surgeons (AAOS).

In the study, 1,657 patients (13 percent of the study population) used [antidepressants](#) up to three years before surgery.

The patients were surveyed before and one year after the THR. The investigators found that a patient's mental health status, assessed by the use of antidepressants before surgery, was a significant factor in predicting outcomes, as well as gender (men are more likely to report lower outcomes), advanced age and co-morbidity (other joint diseases or conditions which affect walking).

According to the investigators, a patient's mental health status should be assessed prior to surgery and taken into consideration during post-operative care.

Provided by American Academy of Orthopaedic Surgeons

Citation: Antidepressant use linked with less patient satisfaction after hip replacement (2012, February 8) retrieved 27 April 2024 from <https://medicalxpress.com/news/2012-02-antidepressant-linked-patient-satisfaction-hip.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.