

# Climate-change effects on malaria risk

February 3 2012, By Adele Rackley

---



A new study suggests that climate change, driven by greenhouse-gas emissions and land-use changes, will cause patterns of malaria infection to change over the next 50 years.

The study combined simulations of mosquito biting rates and [malaria transmission](#) rates with a detailed regional [climate](#) model, to predict changes in the pattern of malaria infection across a large area of Africa up to 2050.

The news is good for some and bad for others. The spread of malaria is predicted to decrease over most parts of tropical Africa because of likely increases in [surface temperatures](#) and reduced [rainfall](#). But epidemics are projected to move south in the sub-Saharan Sahel region, and in much of [East Africa transmission rates](#) of the disease are likely to increase.

In particular the study highlighted the sensitivity of upland areas, where a change of just 2°Celsius could put previously malaria-free regions at risk of epidemics.

"This is one of the first studies to use such a high-resolution climate model across such a wide swathe of Africa," says Dr. Andrew Morse from the University of Liverpool, one of the authors of the research, published in *Environmental Health Perspectives*.

Malaria flourishes in warm, humid environments, so surface temperature and rainfall are the two most important natural factors that influence where it will strike and how quickly it's likely to spread.



Malaria parasites are transmitted to humans by mosquitoes.

The combined weather-disease model showed how temperature and rainfall might change under two climate scenarios using different levels of greenhouse-gas emissions, and also with likely changes to land-use – for example when forests are cleared to make way for crops.

The human effects on climate through land-use change are ignored by many climate models. But they can have a significant effect on local climate – clearing land of natural vegetation, for example, causes local

temperature to rise and reduces rainfall.

The model predicts changes in both the geographic occurrence of malaria and the intensity of transmission.

"Where malaria occurs every year it's already a public health issue," explains Morse. "But we also interested in areas that might be newly at risk of epidemics."

When a malaria epidemic strikes it is in an area where people have low immunity and the disease is more likely to affect people of all ages. So knowing what areas might become at risk in future would make prevention and treatment more effective.

Though it's unrealistic to target malaria treatment programmes as far ahead as 2050, these long-term predictions would be useful for strategic land-use planning, for example where governments are considering developing particular areas or where populations are shifting from rural to urban centres.

One of the benefits of modelling trends at such a small scale is that the researchers can start to narrow down the timescales within which malaria infection patterns could change in particular areas, rather than making general statements. So the next step is to identify which altitude ranges or lowland areas are likely to be affected in a particular ten-year window – information which will be much easier for decision-makers to work with.

The researchers emphasise that other variables not included in this study, like improved prevention and treatment, might in practice counteract the effects of climate on [malaria infection](#). Indirect effects of [climate change](#) on malaria might be important too; for example the effects of a warming climate on agriculture might cause people to migrate to urban

areas, and this would affect transmission rates.

"This model doesn't include human interventions," says Morse, "but having the two climate scenarios and the land-use information shows us how the climate risk may impact on transmission patterns in the future."

**More information:** Ermert V, Fink AH, Morse AP, Paeth H, 2011. The impact of regional climate change on malaria risk due to greenhouse forcing and land-use changes in tropical Africa. *Environ Health Perspect* 120:77-84. doi: 10/1289/ehp.1103681

*This story is republished courtesy of [Planet Earth online](#), a free, companion website to the award-winning magazine Planet Earth published and funded by the Natural Environment Research Council (NERC).*

Provided by PlanetEarth Online

Citation: Climate-change effects on malaria risk (2012, February 3) retrieved 27 April 2024 from <https://medicalxpress.com/news/2012-02-climate-change-effects-malaria.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------