

Preventing and treating drug use with smartphones

February 21 2012

Clinical researchers at the University of Massachusetts Medical School (UMMS) are combining an innovative constellation of technologies such as artificial intelligence, smartphone programming, biosensors and wireless connectivity to develop a device designed to detect physiological stressors associated with drug cravings and respond with user-tailored behavioral interventions that prevent substance use. Preliminary data about the multi-media device, called iHeal, was published online first in the *Journal of Medical Toxicology*.

According to the study's authors, many [behavioral interventions](#) used to treat patients are ineffective outside of the controlled [clinical settings](#) where they are taught. This failure can be attributed to several factors, including a patient's inability to recognize [biological changes](#) that indicate increased risk of relapse and an inability to change their behaviors to reduce health risk.

Edward Boyer, MD, PhD, professor of emergency medicine at UMass Medical School and lead author of the study, worked with colleagues at UMMS and at the Massachusetts Institute of Technology, to design a mobile device using so-called "enabling technologies" that could be used to make behavioral interventions for substance abusers more effective outside the clinic or office environments. The result of their work, iHeal, combines sensors to measure physiological changes and detect trigger points for risky [health behaviors](#), such as substance use, with smartphone software tailored to respond with patient-specific interventions.

Individuals with a history of substance abuse and [post-traumatic stress disorder](#) were asked to wear an iHeal sensor band around their wrist that measures the electrical activity of the skin, body motion, [skin temperature](#) and heart rate – all indicators of stress. The band wirelessly transmits information to a smartphone, where software applications monitor and process the user's physiological data. When the software detects an increased stress level, it asks the user to annotate events by inputting information about their perceived level of stress, drug cravings, and current activities. This information is then used to identify, in real-time, drug cravings and deliver personalized, multimedia drug prevention interventions precisely at the moment of greatest physiological need.

Boyer and his teams examined the iHeal system architecture, as well as preliminary feedback from initial users, to identify key attributes and assess the device's viability. Their analyses suggest a number of technical issues related to data security, as well as the need for a more robust and less stigmatizing version before the device could be worn in public.

Provided by University of Massachusetts Medical School

Citation: Preventing and treating drug use with smartphones (2012, February 21) retrieved 3 May 2024 from <https://medicalxpress.com/news/2012-02-drug-smartphones.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--