

Study quantifies impact of unsafe water and poor sanitation on child and maternal mortality

February 15 2012

The impact of unsafe water and sanitation on the death rates of children under five and mothers in the year after childbirth has been quantified for the first time by Canadian-based researchers.

In a paper published in the UK journal *Environmental Health*, researchers at the United Nations University and McMaster University analyzed data on access to safe water and adequate sanitation across 193 countries.

Using [regression analysis](#) techniques to factor out other variables like income and average children per mother, they compared safe water and sanitation rates with maternal and [child deaths](#) in those countries.

Dividing the 193 countries into four tiers ("quartiles"), they found that countries ranked in the bottom 25% in terms of safe water had about 4.7 more deaths per 1,000 children under five years old compared to countries in the top 25% tier.

The researchers estimate that, when related to safe water access, the difference in mortality between each 25% tier is 1.17 deaths per 1,000 children under five.

Similarly, when judged on access to adequate sanitation, countries ranked in the bottom 25% tier had about 6.6 more deaths per 1,000

children under five years old compared to countries in the top 25% tier.

Put another way, with respect to the availability of [adequate sanitation](#), the difference in mortality between each of the four tiers of countries is estimated at 1.66 deaths per 1,000 children under five.

Relating safe water provision and [maternal death](#) rates (death within a year of [childbirth](#)), the paper says the odds of dying increase 42% from the top tier to each lower tier of countries; the corresponding odds with respect to inadequate sanitation: 48%.

"If the world is to seriously address the [Millennium Development Goals](#) of reducing child and [maternal mortality](#), then improved water and sanitation accesses are key strategies," say Hamilton Canada-based authors June J. Cheng, Susan Watt and Bruce K. Newbold of McMaster University, Andrew Mente of the Population Health Research Institute, and Corinne J. Schuster-Wallace of UNU's Institute for Water, Environment and Health, which initiated the study.

Worldwide, an estimated 1.4 million children die each year from preventable diarrheal diseases and some 88% of diarrhea cases are related to unsafe water, inadequate sanitation or inadequate hygiene.

Earlier World Health Organization (WHO) reports have stated that almost 10% of the global disease burden could be prevented by improving water supply, sanitation, hygiene and management of water resources. Another estimate reports that 4.0% of all deaths and 5.7% of total disability-adjusted life years can be attributed to water, sanitation and hygiene.

Despite progress, previous UN research has shown that, at the current pace, the world will miss the Millennium Development Goal (MDG) for sanitation by 13%, with 2.7 billion people still lacking basic sanitation by

2015. Even if the target was met, 1.7 billion people would still lack access to improved [sanitation](#).

More positively, the world will meet the water MDG target at the current rate. However, even if that forecast is realized by 2015, some 672 million people will still lack access to safe water.

Provided by United Nations University

Citation: Study quantifies impact of unsafe water and poor sanitation on child and maternal mortality (2012, February 15) retrieved 16 April 2024 from <https://medicalxpress.com/news/2012-02-quantifies-impact-unsafe-poor-sanitation.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
