

Snapshot of dialysis: Who's getting treated at home?

February 2 2012

Home-based dialysis treatments are on the rise in both the developing and developed worlds, but developed countries appear to be turning to them less often, according to a study appearing in an upcoming issue of the *Journal of the American Society Nephrology* (JASN). The findings, which provide a global snapshot of dialysis use among patients with kidney failure, may impact future business and research innovations.

Dialysis treatments for [kidney failure](#) come in two forms: peritoneal dialysis is usually done at home, while hemodialysis is predominantly clinic-based. During peritoneal dialysis, fluids that are pumped into the [abdominal cavity](#) collect wastes pulled in from the blood and are then removed from the body. During hemodialysis, blood is removed, purified by a machine, and returned to the body.

Peritoneal dialysis is less expensive than [hemodialysis](#), but its use seems to be declining, particularly in [developed countries](#). To get a sense of the recent trends in peritoneal dialysis use around the world, Arsh Jain, MD (University of Western Ontario, in London, Canada) and his colleagues analyzed records from 1997 to 2008 in 130 countries.

Among the major findings:

- In 2008, there were approximately 196,000 peritoneal dialysis patients worldwide, representing 11% of the dialysis population.
- In total, 59% of peritoneal dialysis patients were treated in

- [developing countries](#) and 41% in developed countries.
- Over the 12 years of the study, the number of peritoneal dialysis patients increased in developing countries by 24.9 patients per million population and in developed countries by 21.8 per million population.
 - The proportion of all dialysis patients treated with peritoneal dialysis did not change in developing countries but significantly declined in developed countries by 5.3%.

These findings indicate that the number of patients treated with peritoneal dialysis rose worldwide from 1997 to 2008, with a 2.5-fold increase in the prevalence of peritoneal dialysis patients in developing countries; however, the proportion of all dialysis patients using this type of treatment continues to decline in developed countries.

"Our findings may impact future business and research innovations," said Dr. Jain. Because of the increasing use of peritoneal dialysis in the developing world and the lower cost of doing research there, "it may be the developing world and not the developed world that drives future medical innovations in peritoneal dialysis," he said.

More information: The article, entitled "Global Trends in Rates of Peritoneal Dialysis," will appear online at on February 2, 2012, [doi:10.1681/ASN.2011060607](https://doi.org/10.1681/ASN.2011060607)

Provided by American Society of Nephrology

Citation: Snapshot of dialysis: Who's getting treated at home? (2012, February 2) retrieved 1 May 2024 from <https://medicalxpress.com/news/2012-02-snapshot-dialysis-home.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.