

# Stimulant treatment for ADHD not associated with increased risk of cardiac events in youth

February 8 2012

---

Attention-deficit/hyperactivity disorder (ADHD) affects 5-9% of youth and is frequently treated with stimulant medications, such as methylphenidate and amphetamine products. A recent safety communication from the US Food and Drug Administration advised that all patients undergoing ADHD treatment be monitored for changes in heart rate or blood pressure.

Amidst growing concern over the risks of stimulant use in youth, a study by Dr. Mark Olfson of the New York State Psychiatric Institute and Columbia University, and his colleagues, published in the February 2012 issue of the *Journal of the American Academy of Child and Adolescent Psychiatry*, assessed the risk of adverse cardiovascular events in children and adolescents without known heart conditions treated with stimulants for ADHD. It is one of the largest studies to date focusing primarily on youth while controlling for pre-existing [cardiovascular risk factors](#).

As reported in the study, Olfson and colleagues examined claims records from a large privately insured population for associations between cardiovascular events in youth with ADHD and stimulant treatment. In total 171,126 privately insured youth aged 6-21 years without known pre-existing heart-related risk factors were followed throughout the study.

The study included patients who have previously received stimulant treatment, patients currently receiving stimulant treatment, and patients

who began or ceased stimulant treatments during the study period. Olfson and colleagues assessed the various groups for incidents of severe cardiovascular events such as [acute myocardial infarction](#), less severe cardiovascular events such as cardiac dysrhythmias, and cardiovascular symptoms such as tachycardia and palpitations. Analysis showed that cardiovascular events and symptoms were rare in this cohort and not associated with stimulant use.

This finding helps to allay concerns of adverse events in otherwise healthy young people receiving treatment for ADHD. Olfson and colleagues said of the results, "It is reassuring that in these young people, short-term stimulant treatment did not substantially increase the risk of [cardiovascular events](#) or symptoms."

**More information:** The article, "Stimulants and Cardiovascular Events in Youth With Attention-Deficit/Hyperactivity Disorder" by Mark Olfson, Cecilia Huang, Tobias Gerhard, Almut G. Winterstein, Stephen Crystal, Paul D. Allison, Steven C. Marcus ([doi:10.1016/j.jaac.2011.11.008](https://doi.org/10.1016/j.jaac.2011.11.008)) appears in *Journal of the American Academy of Child and Adolescent Psychiatry*, Volume 51, Issue 2 (February 2012)

Provided by Elsevier

Citation: Stimulant treatment for ADHD not associated with increased risk of cardiac events in youth (2012, February 8) retrieved 25 April 2024 from <https://medicalxpress.com/news/2012-02-treatment-adhd-cardiac-events-youth.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--