

Health groups issue cervical cancer screening guidelines

March 14 2012

The American Cancer Society (ACS), the American Society for Colposcopy and Cervical Pathology (ASCCP), and the American Society for Clinical Pathology (ASCP) have released new guidelines for the prevention and early detection of cervical cancer. The guidelines generally advise a reduction in the number of tests women get over their lifetime to better ensure that they receive the benefits of testing while minimizing the harms, and include a preference for co-testing using the Pap test and HPV test for women age ages 30 to 65.

The finalized updated [guidelines](#) recommend:

- Women should not be screened before age 21
- Women 21 to 29 should be screened with the [Pap test](#) alone (conventional or liquid-based) every three years. HPV testing should NOT be used for screening in this age group.
- For women 30 and over, the preferred approach is the Pap test plus HPV testing ("co-testing") every five years. Continued screening with the Pap test alone (without HPV testing) every three years is an acceptable alternative. While screening with HPV testing alone is promising, at this time it is not recommended for most [clinical settings](#).
- Screening is not recommended for women over age 65 who have had at least three consecutive negative Pap tests or at least two negative HPV tests the last 10 years, with the most recent test in the last 5 years. Women in this age group who have a history of

pre-cancer (CIN2 or a more severe diagnosis) should continue [routine screening](#) for at least 20 years.

- Women who have undergone a [hysterectomy](#) (with removal of the [cervix](#)) for reasons not related to [cervical cancer](#) or pre-cancer should not be screened.
- Women who have been vaccinated against HPV should follow the age-specific recommendations in these guidelines (for unvaccinated women).

The new guidelines are not intended for women with a history of cervical cancer, exposure to DES in utero, or women who are immunosuppressed (e.g. HIV positive). Costs and other financial issues were not considered in creating the guidelines.

"Pap tests have been done yearly in the past, but we now know that annual screening is not needed and in fact can lead to harm from treatment of cell changes that would never go on to cause cancer," said Debbie Saslow, PhD, director of breast and gynecologic cancer for the [American Cancer Society](#). "Since 1980, organizations including the ACS have recommended less frequent screening. With the addition of the HPV test, we can test even less frequently, as the risk of pre-cancer and cancer when both tests are negative is so low. With these recommendations, our groups are helping to make sure women get the full lifesaving benefits of screening while minimizing its known harms."

Other new recommendations included in the guideline:

- Women at any age should NOT be screened annually by any screening method.
- Women with a slightly abnormal Pap test result (called "ASC-US") and a negative HPV test can be screened again with co-testing in 5 years or with the Pap test alone in 3 years.

- Women with a negative Pap result but a positive HPV test can either be rescreened with co-testing in one year, or tested with a test for specific types of HPV (HPV16 and HPV 18).

The updated guidelines were first released in draft form in late 2011. The working groups that created the draft guidelines then met with delegates from 25 organizations to further discuss and finalize the recommendations, which were then adapted into this final guideline.

"Our process resulted in guidelines that are focused on collectively presenting the best patient-centered cervical cancer screening strategies," said Mark Stoler, MD, past-president of the American Society for Clinical Pathology. "These final recommendations are based on a broad and emerging body of literature, and meld the very latest knowledge on the interplay between new molecular tests and traditional cytology."

"While these new guidelines reflect relatively small changes over previous [screening](#) recommendations, they are important," said Alan Waxman, M.D., incoming president of the American Society for [Colposcopy](#) and Cervical Pathology. "The addition of HPV testing to the Pap test in [women](#) 30 and over has been shown in recent studies to provide better protection for longer intervals from cancer and pre-cancerous changes than the use of the Pap test alone."

The guidelines are being published jointly in *CA: A Cancer Journal for Clinicians* (ACS), *Journal of Lower Genital Tract Disease* (ASCCP), and *American Journal of Clinical Pathology* (ASCP).

More information: "Screening Guidelines for the Prevention and Early Detection of Cervical Cancer," Saslow et al. *CA Cancer J Clin* 2012;00:000-000; [doi:10.3322/caac.21139](https://doi.org/10.3322/caac.21139)

Provided by American Cancer Society

Citation: Health groups issue cervical cancer screening guidelines (2012, March 14) retrieved 4 May 2024 from <https://medicalxpress.com/news/2012-03-health-groups-issue-cervical-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.