

# Medical researchers recommend male circumcision

March 2 2012

---

Penile cancer, HIV, HPV, syphilis and kidney inflammation are among a number of medical conditions whose risk can be lowered by the practice of infant male circumcision, says a study led by the University of Sydney.

The review of current international evidence for the benefits and risks of infant male circumcision is published today in the [Open Journal of Preventive Medicine](#).

"This is the world's first evidence-based policy on infant circumcision and its authors include five Fellows of the Royal Australasian College of Physicians," lead author, Professor Brian Morris, from Sydney Medical School at the University, said.

"The evidence in favor of infant circumcision is now so strong that advocating this simple, inexpensive procedure for baby boys is about as effective and safe as childhood vaccination".

The study points out that many common childhood conditions, including [kidney damage](#), will become rare if baby boys are circumcised in the first weeks of life.

Circumcision was found to protect men, and their sexual partners, from several common sexually transmitted infections, as well as cancers of the penis and cervix.

"The scientific evidence shows no adverse effects on sexual function, sensitivity, satisfaction or sensation - if anything the opposite," said Professor Morris.

The policy suggests that for maximum benefits, safety, convenience and cost savings, circumcision should be performed in infancy and with local anaesthesia. It claims that the risk-benefit analysis shows benefits outweigh minor risks by a factor of over 100 to 1.

"It is now up to state governments to ensure that bans on elective infant [male circumcision](#) in [public hospitals](#) are lifted without delay. And it is essential that the federal government revises the Medicare rebate so that this procedure is affordable for low-income families.

"The costs saved will be enormous, as this policy statement shows that half of uncircumcised males will suffer an adverse medical condition over their lifetime, and many will die as a result of diseases preventable by circumcision," Professor Morris said.

Provided by University of Sydney

Citation: Medical researchers recommend male circumcision (2012, March 2) retrieved 20 March 2024 from <https://medicalxpress.com/news/2012-03-medical-male-circumcision.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------