

Success in growing functional pituitary gland in a lab culture will advance regenerative medicine

March 2 2012



Figure 1: A micrograph of an embryonic pituitary gland that self-formed in an embryonic stem cell aggregate on day 13 of culture. Credit: Reproduced from Ref. 1 © 2012 Yoshiki Sasai

Embryonic stem cells grown in a laboratory culture can organize themselves into a partial pituitary gland that is fully functional when transplanted into mice, a team of researchers led by Yoshiki Sasai of the RIKEN Center for Developmental Biology reports. These researchers developed a novel cell-culture technique for growing stem cells in threedimensional floating clusters. They had previously shown that stem cells



grown in this way can organize themselves into functional eye and brain tissue.

In their study on the <u>pituitary gland</u>, the researchers used their technique to grow mouse stem cells, and then altered the culture conditions to recapitulate the embryonic environment that gives rise to the anterior pituitary, or adenohypophysis.

The adenohypophysis contains cells that synthesize and secrete five hormones under the control of the hypothalamus. After release into the blood stream, these hormones play multiple roles in the body, including regulation of growth, blood pressure, metabolism and sex-organ function. Development of the adenohypophysis requires interaction between two types of tissue, leading to the formation of a small pouch that pinches off from the area of the embryo and forms the mucosa in the mouth.

When the researchers stimulated stem-cell clusters with specific signaling molecules, they generated both tissue types that separated naturally into layers. Cells at the interface between the two layers then spontaneously formed oval-shaped pouch-like structures (Fig. 1), before differentiating into four types of precursors, each of which began to synthesize and secrete a different hormone.

Sasai and colleagues transplanted the clusters into the kidneys of mice whose pituitaries had been surgically removed. Normally these mice would die two months post-surgery, but the transplanted cells rescued the animals by normalizing the levels of hormones in the bloodstream.

The research could open new avenues for stem-cell therapies to treat hormonal disorders. It also represents a significant advance in using <u>stem</u> <u>cells</u> to generate complex three-dimensional structures: it represents a step towards growing fully functional organs in the laboratory.



This technique has a straightforward medical application in treating human patients with pituitary deficiencies. "Growth hormone deficiency could be also targets of the cell therapy," Sasai notes. Eventually, it could be possible to grow functional neural tissue for transplantation into the human brain. For example, the ability to grow cerebellar tissue containing Purkinje cells could be useful for treating cerebellar atrophy.

"Regenerative medicine is surely proceeding in this direction," says Sasai. "We are now developing computer-based models and simulations to facilitate the design of more complex organs."

More information: Suga, H., et al. Self-formation of functional adenohypophysis in three-dimensional culture. <u>Nature</u> 480, 57–62 (2011).

Eiraku, M., et al. Self-organizing optic-cup morphogenesis in threedimensional culture. <u>Nature</u> 472, 51–56 (2011).

Eiraku, M., et al. Self-organized formation of polarized cortical tissues from ESCs and its active manipulation by extrinsic signals. <u>Cell Stem</u> <u>Cell</u> 3, 519–532 (2008).

Provided by RIKEN

Citation: Success in growing functional pituitary gland in a lab culture will advance regenerative medicine (2012, March 2) retrieved 16 June 2024 from <u>https://medicalxpress.com/news/2012-03-success-functional-pituitary-gland-lab.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.