

Study may offer clues to reverse cognitive deficits in humans

April 26 2012

The ability to navigate using spatial cues was impaired in mice whose brains were minus a channel that delivers potassium — a finding that may have implications for humans with damage to the hippocampus, a brain structure critical to memory and learning, according to a Baylor University researcher.

Mice missing the channel also showed diminished learning ability in an experiment dealing with fear conditioning, said Joaquin Lugo, Ph.D., the lead author in the study and an assistant professor of psychology and neuroscience in Baylor's College of Arts & Sciences.

"By targeting chemical pathways that alter those [potassium](#) channels, we may eventually be able to apply the findings to humans and reverse some of the cognitive deficits in people with epilepsy and other neurological disorders," Lugo said.

The research was done in Baylor College of Medicine Intellectual and Developmental Disabilities Research Center Mouse Neurobehavior Core in Houston during Lugo's time as a researcher there.

The findings are published online in the journal *Learning & Memory*.

The channel, called Kv4.2, delivers potassium, which aids neuron function in the brain's hippocampus. The [hippocampus](#) forms memory for long-term storage in the [brain](#). Potassium also helps to regulate excitability.

Individuals who have epilepsy sometimes exhibit altered or missing Kv4.2 channels or similar types of channels.

In the experiment investigating navigation, "knockout" mice — those without the channel — were tested in a water maze four feet in diameter and 12 inches deep, with eight trials daily — each lasting about a minute — over four days, he said. Their performance was compared with that of normal mice.

Both groups responded to visual cues — colored symbols — in learning their way around the maze, but the knockout mice did not respond as well as the normal mice in terms of spatial cues — hidden platforms in the water.

"When the mice don't have this channel, it hurts their ability to learn," Lugo said.

In a separate experiment examining fear conditioning, both [knockout mice](#) and normal mice were placed in a cage, and researchers sounded a tone before giving the mice a mild electric shock. In repeated trials, both groups began to freeze upon hearing the tone as they anticipated a shock. But the normal mice also reacted to the context — being placed in the cage — while the mice who did not have the Kv4.2 channel reacted only to the tone.

More information: <http://learnmem.cshlp.org/content/19/5/182.full>

Provided by Baylor University

Citation: Study may offer clues to reverse cognitive deficits in humans (2012, April 26) retrieved 30 April 2024 from

<https://medicalxpress.com/news/2012-04-clues-reverse-cognitive-deficits-humans.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.