


Complex role of genes in autism revealed

April 4 2012



At the University of Washington in Seattle, Dr. Brian O'Roak studies genetic origins of intellectual, social and psychiatry disabilities in children. Credit: Clare McLean

University of Washington researchers announced their findings from a major study looking into the genetic basis of autism spectrum disorders (ASD) with an approach piloted at the UW. Their results are reported in the April 4 advanced online edition of the journal *Nature*.

The researchers have been studying ASD in children who have no family history of this or related impairments — so called "sporadic autism" — and also why autism varies in its symptoms and severity. By focusing on "sporadic autism," the researchers sought to evaluate a specific genetic model for ASD risk, namely the appearance of new mutations (termed *de novo*) in children with ASD that were not found in either parent.

By uncovering new gene mutations that disrupt the function of proteins, the researchers have discovered a pathway related to modifying chromatin — the tightly coiled spools of DNA in the cell — and to regulating genes in the brain and nervous system. Various changes in this pathway contribute to children developing autism in different ways. Mutations in this pathway also may contribute to a variety of childhood intellectual, social, and psychiatric disabilities, with implications beyond autism.

To identify these new mutations, the researchers used the latest sequencing technologies and analytical methods to determine the sequence of the protein-coding portion of the human genome, called the "exome," in family trios (father, mother, and child). This approach was piloted this past year at the University of Washington with an initial set of 20 autism families. The pilot demonstrated the technical feasibility and potential impact of this approach.

For the current study, the researchers expanded the project to include 677 individuals from 209 families with a single child with autism. They also sequenced the exomes of 50 unaffected brothers and sisters. In the newly reported results, 248 de novo mutations were validated, and 120 of these were classified as severe. These were predicted to produce, for example, proteins that were truncated or malfunctioning. The researchers then narrowed in on 60 top candidates most likely to contribute to autism risk, based on the nature of the mutation, functional evidence, or previous studies.

"It is important to point out that in each generation there is on average one new coding mutation per child and not all of these will cause developmental problems. However, in the case of children with autism, what we are finding are disruptions in many genes that are known to directly interact and also look similar to genes previously associated with autism," said Dr. Brian J. O'Roak, a senior fellow in the Department of

Genome Sciences working with senior authors Drs. Jay Shendure and Evan Eichler. In fact, researchers found that 49 of the genes mutated had products known to directly interact by forming a highly interconnected network.

Interestingly, many of the proteins in this pathway are important in terms of remodeling chromatin — changing the way DNA is packaged in the cell —and controlling the expression and function of other genes and proteins. These protein pathways are thought to be critical in brain cell formation, brain cell connections, and nerve-cell signaling.

Having this large data set also allowed the researchers to evaluate the parental source of these new mutations, that is, whether they came from the sperm of the father or egg of the mother. Their analysis revealed that the new mutations were overwhelming paternal in origin (in a ratio of 4:1). Their results confirmed a prediction population geneticist J.B.S. Haldane made in in 1935. Moreover, the new mutations occurred at a rate that correlated with the age of the father. These findings, they said, support other studies that show older fathers have a slightly increased risk of having a child with an autism spectrum disorder.

What is also very clear from this study and two additional studies appearing concurrently in the same issue of *Nature* is that autism risk mutations are scattered across many genes. One of the other studies was led by Mount Sinai Medical Center in New York, the second by Yale University in New Haven.

In the UW study, recurrent protein-altering mutations were discovered in only two genes, NTNG1, and CHD8. The data suggest that, at the molecular level, there are many different forms of autism and that the term "autism spectrum disorder" is better thought of as an umbrella disorder with many root causes. The authors predict that although no single gene will account for more than 1 percent of autism, collectively

all of these rare mutations will account for much of the genetic basis of the disease.

While this level of complexity is a major challenge for the field, the authors are already working on a solution using next-generation sequencing approaches. To tackle this challenge, the researchers implemented a new cost-effective screening technology that allowed them to screen more than 2,500 individuals for mutations in six genes in only a month. In so doing they found strong evidence for the involvement of a glutamate receptor gene GRIN2B for a subset of cases with autism. This same approach will allow the authors to screen all of the newly discovered genes to rapidly test which ones are truly disease-causing.

Among the other genes they discovered with de novo mutations in children with autism, several have been previously implicated in intellectual disability and developmental delay. This indicates, the authors said, that the divisions clinicians made between these various types of diseases in children may not readily translate into differences at the molecular level. The researchers added that it is still uncertain whether there are subsets of people with autism who share a common or strongly related causative mechanism in their underlying molecular biology, or how large those groups might be.

More information: Neale et al. Patterns and rates of exonic de novo mutations in autism spectrum disorders. *Nature* DOI: 10.1038/nature11011

Provided by University of Washington

Citation: Complex role of genes in autism revealed (2012, April 4) retrieved 4 May 2024 from

<https://medicalxpress.com/news/2012-04-complex-role-genes-autism-revealed.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.