

One in three PIP implants may rupture: British surgeons

April 17 2012

A pair of British plastic surgeons said Tuesday the rupture rate of PIP breast implants behind a global health scare may be as high as one in three, significantly more than previously thought.

A study of 453 of their patients who had received PIP implants between seven and 12 years ago showed a rupture rate of between 15.9 percent and 33.8 percent.

"Previous studies have typically reported failure rates of between two percent and five percent," said authors Miles Berry and Jan Stanek of the study published in the Journal of Plastic, Reconstructive and [Aesthetic Surgery](#).

The new study had employed more conclusive ultrasound scans rather than the physical examinations typically used, they said.

"All PIP implants, due to the high rupture rate and uncertainty about the nature of the silicone gel, may need to be removed," Stanek was quoted as saying in a statement, adding that further research must be done.

"Being only a single-handed practice, this initial study is the tip of an iceberg that may affect 40,000 [women](#) in the UK with PIP implants," added a statement.

More than 400,000 women around the world are believed to have received implants made by French company PIP, which shut in 2010

after it was found to have used substandard, industrial-grade silicone gel.

Of them, about 47,000 were in Britain, 30,000 in France and 4,300 in Italy.

France, Germany, the Czech Republic and the Netherlands have recommended the devices be removed as a precaution.

Britain has not followed suit, but has said that women who had their implants on the [National Health Service](#) (NHS) could have them replaced without charge.

More than a dozen countries in Europe and Latin America have urged women to seek regular checkups.

Manufacturers and plastic surgeons say that [breast implants](#) have an expected, normal lifespan of about 12 years.

Usually in the case of rupture the [silicone gel](#) does not leak, but even if it does should not cause tissue irritation. However, in the case of the PIP implants, the gel used was of a non-medical grade.

"Those patients with ruptured implants will have to have them removed; those with no evidence of rupture will need to be monitored on a regular basis," said Stanek.

"Further research into the nature of the elastomer (outer casing) and [gel](#) filler will determine whether all [PIP implants](#) should be explanted in the future."

(c) 2012 AFP

Citation: One in three PIP implants may rupture: British surgeons (2012, April 17) retrieved 6

May 2024 from <https://medicalxpress.com/news/2012-04-pip-implants-rupture-british-surgeons.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.