


Uric acid levels predict death in acute coronary syndrome

April 20 2012



Elevated uric acid levels are predictive of one-year mortality in patients with acute coronary syndrome, according to a study published in the May 1 issue of *The American Journal of Cardiology*.

(HealthDay) -- Elevated uric acid levels are predictive of one-year mortality in patients with acute coronary syndrome, according to a study published in the May 1 issue of *The American Journal of Cardiology*.

Gjin Ndrepepa, M.D., of the Deutsches Herzzentrum in Munich, Germany, and colleagues conducted a study involving 5,124 patients with [acute coronary syndrome](#) who underwent percutaneous coronary intervention (PCI) to evaluate whether uric acid levels have any prognostic value in this patient population. Of the participants, 1,629 had acute ST-segment elevation [myocardial infarction](#) (STEMI), 1,332 had acute non-STEMI, and 2,163 had [unstable angina](#). Participants were

classified into four uric acid quartiles: quartile 1 (1.3 to

Citation: Uric acid levels predict death in acute coronary syndrome (2012, April 20) retrieved 24 April 2024 from <https://medicalxpress.com/news/2012-04-uric-acid-death-acute-coronary.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.