

Seeing Beyond the Visual Cortex

April 3 2012, By Miles O' Brien and Jon Baime

(Medical Xpress) -- It's a chilling thought--losing the sense of sight because of severe injury or damage to the brain's visual cortex. But, is it possible to train a damaged or injured brain to "see" again after such a catastrophic injury? Yes, according to Tony Ro, a neuroscientist at the City College of New York, who is artificially recreating a condition called blindsight in his lab.

"Blindsight is a condition that some patients experience after having damage to the [primary visual cortex](#) in the back of their brains. What happens in these patients is they go cortically blind, yet they can still discriminate visual information, albeit without any awareness." explains Ro.

While no one is ever going to say blindsight is 20/20, Ro says it holds tantalizing clues to the architecture of the brain. "There are a lot of areas in the brain that are involved with processing [visual information](#), but without any [visual awareness](#)," he points out. "These other parts of the brain receive input from the eyes, but they're not allowing us to access it consciously."

With support from the National Science Foundation's (NSF) Directorate for Social, Behavioral and Economic Sciences, Ro is developing a clearer picture of how other [parts of the brain](#), besides the visual cortex, respond to [visual stimuli](#).

In order to recreate blindsight, Ro must find a volunteer who is willing to temporarily be blinded by having a powerful magnetic pulse shot right

into their visual cortex. The magnetic blast disables the visual cortex and blinds the person for a split second. "That blindness occurs very shortly and very rapidly--on the order of one twentieth of a second or so," says Ro.

On the day of Science Nation's visit to Ro's lab in the Hamilton Heights section of Manhattan, volunteer Lei Ai is seated in a small booth in front of a computer with instructions to keep his eyes on the screen. A round device is placed on the back of Ai's head. Then, the booth is filled with the sound of consistent clicks, about two seconds apart. Each click is a magnetic pulse disrupting the activity in his [visual cortex](#), blinding him. Just as the pulse blinds him, a shape, such as a diamond or a square, flashes onto a computer screen in front of him.

Ro says that 60 to nearly 100 percent of the time, test subjects report back the shape correctly. "They'll be significantly above chance levels at discriminating those shapes, even though they're unaware of them. Sometimes they're nearly perfect at it," he adds.

Ro observes what happens to other areas of Ai's brain during the instant he is blinded and a shape is flashed on the screen. While the [blindness](#) wears off immediately with no lasting effects, according to Ro, the findings are telling. "There are likely to be a lot of alternative visual pathways that go into the brain from our eyes that process information at unconscious levels," he says.

Ro believes understanding and mapping those alternative pathways might be the key to new rehabilitative therapies. "We have a lot of soldiers returning home who have a lot of brain damage to visual areas of the [brain](#). We might be able to rehabilitate these patients," he says. And that's something worth looking into.

Provided by National Science Foundation

Citation: Seeing Beyond the Visual Cortex (2012, April 3) retrieved 28 April 2024 from <https://medicalxpress.com/news/2012-04-visual-cortex.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.