

Cardiac disease risk factors prevalent among U.S. teens

May 21 2012



(HealthDay) -- From 1999 to 2008 the prevalence of several cardiovascular disease risk factors remained stable among U.S. adolescents, but the burden of risk factors is still considerable, according to a study published online May 21 in *Pediatrics*.

To investigate recent trends in the prevalence of selected CVD risk factors among U.S. <u>adolescents</u>, Ashleigh L. May, Ph.D., from the U.S. <u>Centers for Disease Control and Prevention</u> in Atlanta, and colleagues analyzed data from 3,383 12- to 19-year-old participants from the 1999 to 2008 National Health and <u>Nutrition Examination Survey</u>.

The researchers found that from 1999 to 2008 the overall prevalence of prehypertension/hypertension was 14 percent, borderline-high/high low-density lipoprotein (LDL) cholesterol was 22 percent, low high-density



lipoprotein cholesterol (

Citation: Cardiac disease risk factors prevalent among U.S. teens (2012, May 21) retrieved 9 April 2024 from

https://medicalxpress.com/news/2012-05-cardiac-disease-factors-prevalent-teens.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.