

Commonly used painkillers may protect against skin cancer

May 29 2012

A new study suggests that aspirin and other similar painkillers may help protect against skin cancer. Published early online in *Cancer*, a peer-reviewed journal of the American Cancer Society, the findings indicate that skin cancer prevention may be added to the benefits of these commonly used medications.

Previous studies suggest that taking nonsteroidal anti-inflammatory drugs, or NSAIDs, which include aspirin, ibuprofen, and naproxen, as well as a variety of other nonprescription and [prescription drugs](#), can decrease an individual's risk of developing some [types of cancer](#). Sigrún Alba Jóhannesdóttir, BSc, of Aarhus University Hospital in Denmark, and her colleagues looked to see if the medications might decrease the risk of the three major types of skin cancer: basal cell carcinoma, squamous cell carcinoma, and malignant melanoma.

The researchers analyzed medical records from northern Denmark from 1991 through 2009 and identified 1,974 diagnoses of squamous cell carcinoma, 13,316 diagnoses of basal cell carcinoma, and 3,242 diagnoses of malignant melanoma. They compared information, including prescription data, from these patients with information from 178,655 individuals without skin cancer.

Individuals who filled more than two prescriptions for NSAIDs had a 15 percent decreased risk for developing squamous cell carcinoma and a 13 percent decreased risk for developing malignant melanoma than those who filled two or fewer prescriptions for the medications, especially

when the drugs were taken for seven or more years or taken at high intensity. Individuals who took NSAIDs did not seem to benefit from a reduced risk of developing basal cell carcinoma in general, although they did have a 15 percent and 21 percent reduced risk of developing this type of cancer on less-exposed sites (body areas other than the head and neck) when they took them long term or at high intensity, respectively.

"We hope that the potential cancer-protective effect of [NSAIDs](#) will inspire more research on [skin cancer](#) prevention," said Ms. Jóhannesdóttir. "Also, this potential cancer-protective effect should be taken into account when discussing benefits and harms of NSAID use."

More information: "Nonsteroidal anti-inflammatory drugs and the risk of skin cancer: A population-based case-control study." Sigrún Alba Jóhannesdóttir, Ellen T. Chang, Frank Mehnert, Morten Schmidt, Anne Braae Olesen, Henrik Toft Sørensen. *CANCER*; Published Online: May 29, 2012 ([DOI: 10.1002/cncr.27406](https://doi.org/10.1002/cncr.27406)).

Provided by Wiley

Citation: Commonly used painkillers may protect against skin cancer (2012, May 29) retrieved 4 May 2024 from <https://medicalxpress.com/news/2012-05-commonly-painkillers-skin-cancer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
