

Highly targeted irradiation as good as whole breast radiotherapy in early stage cancer

May 11 2012

Barcelona, Spain: Using a concentrated, highly targeted dose of radiation to the breast has equally good results as irradiating the whole area, with no adverse effects on survival and a much better cosmetic outcome, Hungarian researchers have found. Reporting the ten-year results of a randomised trial, Professor Csaba Polgár, MD, Director of the Centre for Radiotherapy, National Institute of Oncology, Budapest, will tell the World Congress of Brachytherapy [1 & 2] today (Friday) that he believes that accelerated partial breast irradiation (APBI) could be offered to many more breast cancer patients, resulting in fewer side effects and major cost savings to healthcare systems.

The Budapest trial was the first randomised study to compare the efficacy and [side effects](#) of conventional whole [breast irradiation](#) with APBI using multicatheter brachytherapy. This technique consists of inserting multiple plastic catheters into the tumour bed after breast conserving surgery. The catheters are inserted under local or general anaesthesia, according to the wishes of the patient. The radiation source is then loaded into them to deliver a highly concentrated dose to the precise region of the excised tumour – the region most likely to develop a local recurrence of the cancer. Using APBI means that the major part of the healthy breast tissue does not receive radiation.

"In a group of 258 women who had had breast-conserving surgery, we found that, at a median follow-up time of just over ten years, the rate of local recurrence – the cancer coming back in the same place – was 5.9% in the APBI group and 5.1% in the group who received whole breast

irradiation," says Professor Polgár. "There was no significant difference in overall [survival](#), cancer-specific survival, and disease free survival between the two groups. However, the differences in cosmetic outcome were striking – 81% in the APBI group had good-to-excellent results as opposed to 63% of those who received whole breast irradiation."

A further advantage of APBI is that treatment time can be shortened. As opposed to whole breast irradiation, where a course of treatment typically takes five weeks (five treatments per week), APBI can be carried out in four to five days. "We believe that the results of our trial will encourage more patients to choose APBI, particularly those who may now be inclined to refuse radiotherapy because of the logistical difficulties involved – having to travel long distances over a period of six weeks can be a major problem for some patients," Professor Polgár will say.

Fewer than 10% of early stage breast cancer patients currently receive APBI in Europe, and in some countries it is still considered an investigational treatment. In other parts of the world, for example the USA, it has already been accepted into general practice for the treatment of selected low-risk patients.

"Our trial has shown that it is possible to use a faster and more focused treatment without compromising survival. It is a little like comparing a total mastectomy with a lumpectomy – we have found that we can improve cosmetic outcomes, lessen side effects, and achieve greater patient satisfaction with a less invasive and more cost-effective technique. Other multi-centre trials of APBI are currently underway, and we believe that these are likely to give further support to our findings," Professor Polgár will conclude.

Dr. Christine Haie-Meder, [Radiation](#) Oncologist at the Institut Gustave Roussy, Villejuif, France and chairperson of the congress, said: "We

know that all the patients with [breast cancer](#) do not need whole breast irradiation. Especially in elderly patients, the treatment, when limited to partial breast irradiation, can be significantly shortened. This study is important because, within the frame of a [randomised trial](#) with a ten-year follow-up, it shows that in selected patients, partial breast irradiation gives as good results as whole [breast](#) irradiation with the same recurrence rate, but with significantly better cosmetic results."

More information: [1] This year the World Congress of Brachytherapy is held in parallel with the 31st conference of the European Society for Radiotherapy and Oncology (ESTRO 31), both taking place in Barcelona between May 9-13.

[2] Brachytherapy is a type of internal radiotherapy, which involves putting solid radioactive material close to, or inside, the tumour. This gives a high dose of radiotherapy directly to the tumour but only a low dose to normal tissues.

Provided by European Society for Radiotherapy and Oncology

Citation: Highly targeted irradiation as good as whole breast radiotherapy in early stage cancer (2012, May 11) retrieved 27 April 2024 from <https://medicalxpress.com/news/2012-05-highly-irradiation-good-breast-radiotherapy.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--