

# Psychopathy linked to specific structural abnormalities in the brain

May 7 2012

---

New research provides the strongest evidence to date that psychopathy is linked to specific structural abnormalities in the brain. The study, published in *Archives of General Psychiatry* and led by researchers at King's College London is the first to confirm that psychopathy is a distinct neuro-developmental sub-group of anti-social personality disorder (ASPD).

Most [violent crimes](#) are committed by a small group of persistent male offenders with ASPD. Approximately half of male prisoners in England and Wales will meet [diagnostic criteria](#) for ASPD. The majority of such men are not true psychopaths (ASPD-P). They are characterised by emotional instability, [impulsivity](#) and high levels of mood and [anxiety disorders](#). They typically use aggression in a reactive way in response to a perceived threat or sense of frustration.

However, about one third of such men will meet additional diagnostic criteria for psychopathy (ASPD+P). They are characterised by a lack of empathy and remorse, and use aggression in a planned way to secure what they want (status, money etc.). Previous research has shown that psychopaths' brains differ structurally from healthy brains, but until now, none have examined these differences within a population of violent offenders with ASPD.

Dr Nigel Blackwood from the Institute of Psychiatry at King's and lead author of the study says: 'Using [MRI scans](#) we found that psychopaths had structural [brain abnormalities](#) in key areas of their 'social brains'

compared to those who just had ASPD. This adds to behavioural and developmental evidence that psychopathy is an important subgroup of ASPD with a different neurobiological basis and different treatment needs.

'There is a clear behavioural difference amongst those diagnosed with ASPD depending on whether or not they also have psychopathy. We describe those without psychopathy as 'hot-headed' and those with psychopathy as 'cold-hearted'. The 'cold-hearted' psychopathic group begin offending earlier, engage in a broader range and greater density of offending behaviours, and respond less well to treatment programmes in adulthood, compared to the 'hot-headed' group. We now know that this behavioural difference corresponds to very specific structural brain abnormalities which underpin psychopathic behaviour, such as profound deficits in empathising with the distress of others.'

The researchers used Magnetic Resonance Imaging (MRI) to scan the brains of 44 violent adult male offenders diagnosed with Anti-Social Personality Disorder (ASPD). Crimes committed included murder, rape, attempted murder and grievous bodily harm. Of these, 17 met the diagnosis for psychopathy (ASPD+P) and 27 did not (ASPD-P). They also scanned the brains of 22 healthy non-offenders.

The study found that ASPD+P offenders displayed significantly reduced grey matter volumes in the anterior rostral prefrontal cortex and temporal poles compared to ASPD-P offenders and healthy non-offenders. These areas are important in understanding other people's emotions and intentions and are activated when people think about moral behaviour. Damage to these areas is associated with impaired empathising with other people, poor response to fear and distress and a lack of 'self-conscious' emotions such as guilt or embarrassment.

Dr Blackwood explains: 'Identifying and diagnosing this sub-group of

violent offenders with brain scans has important implications for treatment. Those without the syndrome of [psychopathy](#), and the associated structural brain damage, will benefit from cognitive and behavioural treatments. Optimal treatment for the group of psychopaths is much less clear at this stage.'

**More information:** Gregory, S. et al. 'The Antisocial Brain: Psychopathy Matters – a structural MRI investigation of antisocial male offenders', *Archives of General Psychiatry* (7th May 2012)

Provided by King's College London

Citation: Psychopathy linked to specific structural abnormalities in the brain (2012, May 7)  
retrieved 24 April 2024 from  
<https://medicalxpress.com/news/2012-05-psychopathy-linked-specific-abnormalities-brain.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--