

# Simple, low-cost checklist dramatically improves practices of health workers during childbirth

May 16 2012

---

Boston, MA — A new study from Harvard School of Public Health (HSPH) and the World Health Organization (WHO) found that a simple checklist-based childbirth safety program dramatically improved adherence to essential childbirth care practices at a pilot hospital in south India. Of 29 practices measured, 28 were improved after adoption of the checklist and overall adherence to essential practices was 150% better after the checklist was introduced.

"This is a significant step forward because it provides hope that use of this simple, low-cost tool can help birth attendants better adhere to universally accepted standards in [childbirth](#) care," said senior author Atul Gawande, professor in health policy and management at HSPH, lead advisor for WHO Patient Safety's childbirth and surgery safety programs, and a surgeon at Brigham and Women's Hospital, Boston.

The study appears in the May 16, 2012 online edition of *PLoS ONE*.

Nearly 300,000 maternal deaths, 3.1 million newborn deaths, and 1.2 million intrapartum-related stillbirths take place in low-income countries each year; the vast majority are preventable. From 2008 to 2010, HSPH and WHO developed the WHO Safe Childbirth Checklist program to address the major causes of maternal and neonatal mortality. As part of its development, the checklist was field tested for usability in 10 countries, mostly in Africa and south Asia.

The researchers conducted the study at a hospital in Karnataka, India. As part of the checklist-based program, local staff reviewed deficiencies in their current practices and undertook training on using the WHO Safe Childbirth Checklist. Items on the 29-item checklist address the major causes of maternal deaths (e.g. hemorrhage, infection, obstructed labor, and hypertensive disease), intrapartum-related stillbirths, and neonatal deaths (e.g. complications of premature birth, infection, and birth asphyxia).

Researchers observed the childbirth practices of health care workers during 499 birth events—the period from admission to discharge—prior to introducing the checklist to establish a baseline, and then compared the results with 795 birth events after implementing the checklist.

The results reveal that the number of essential practices performed by the hospital workers increased from an average of 10 of 29 at baseline to 25 of 29 after implementing the checklist. "The checklist program actively prompted health care workers to remember to complete proven practices such as handwashing, infection management, postpartum bleeding assessment, and breastfeeding within an hour after birth," said Dr. Bhala Kodkany, professor of obstetrics and gynecology and director of JNMC Women's and Children's Health Research Unit in Karnataka, India, and a co-investigator.

Promoting institutional delivery in order to facilitate skilled care by trained birth attendants is a major global strategy to improve the safety of childbirth. This approach has resulted in increasing numbers of births taking place in already overburdened (and often under-resourced) facilities. Without low-cost, simple and effective methods to improve quality of care in these facilities, health outcomes are not improving at the rate they should be.

"The results of this study are the first evidence to suggest that the

success we've seen with checklists in other health disciplines, for example in surgery, might also be applied to prevent avoidable childbirth-related deaths in low-income countries," said Itziar Larizgoitia, coordinator in the Patient Safety Programme for the World Health Organization.

This study measured the way health workers care for women and newborns during childbirth but was too small a study to measure the impact on complications or reducing deaths. The researchers are now conducting a large-scale trial in more than 100 hospitals in north India to determine if the [checklist](#) program can save the lives of women and newborns.

**More information:** "Improving Quality of Care for Maternal and Newborn Health: Prospective Pilot Study of the WHO Safe Childbirth Checklist Program," Jonathan M. Spector, Priya Agrawal, Bhala Kodkany, Stuart Lipsitz, Angela Lashoher, Gerald Dziekan, Rajiv Bahl, Mario Merialdi, Matthews Mathai, Claire Lemer, Atul A. Gawande, PLoS One, online May 16, 2012

Provided by Harvard School of Public Health

Citation: Simple, low-cost checklist dramatically improves practices of health workers during childbirth (2012, May 16) retrieved 2 May 2024 from <https://medicalxpress.com/news/2012-05-simple-low-cost-checklist-health-workers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
---