

Correct treatment of common diabetic foot infections can reduce amputations

May 22 2012

Diabetic foot infections are an increasingly common problem, but proper care can save limbs and, ultimately, lives, suggest new guidelines released by the Infectious Diseases Society of America (IDSA).

Poor treatment of infected foot wounds in people with diabetes can lead to lower extremity amputation, and about 50 percent of patients who have foot amputations die within five years – a worse mortality rate than for most cancers. But about half of lower extremity amputations that aren't caused by trauma can be prevented through proper care of foot infections, note the new IDSA diabetic foot infections guidelines, which are being published today in the journal *Clinical Infectious Diseases*.

Because people with diabetes often have poor circulation and little or no feeling in their feet, a sore caused by a rubbing shoe or a cut can go unnoticed and worsen. As many as one in four people with diabetes will have a foot ulcer – an open sore – in their lifetime. These wounds can easily become infected. Unchecked, the infection can spread, killing soft tissue and bone. Dead and infected tissue must be surgically removed, which, if the infection is extensive, can mean amputation of the toe, foot, or even part of the leg. Nearly 80 percent of all nontraumatic amputations occur in people with diabetes – and 85 percent of those begin with a foot ulcer.

"Lower extremity amputation takes a terrible toll on the diabetic patient," said Benjamin A. Lipsky, MD, chair of the review panel, lead author of the guidelines and professor of medicine at the University of

Washington and VA Puget Sound, Seattle. "People who have had a foot amputated often can no longer walk, their occupational and social opportunities shrink, and they often become depressed and are at significant risk for a second amputation. Clearly, preventing amputations is vital, and in most cases, possible."

The guidelines emphasize the importance of rapid and appropriate therapy for treating infected wounds on the feet, typically including surgical removal (debridement) of dead tissue, appropriate antibiotic therapy and, if necessary, removing pressure on the wound and improving blood flow to the area. Many patients with foot infections initially receive only antibiotic therapy, which is often insufficient in the absence of proper wound care and surgical interventions, the guidelines note.

Because treatment of diabetic foot infections can be complicated, the best approach is to involve a multidisciplinary team that can assess and address various aspects of the problem, suggest the guidelines, which are a revision and update of IDSA's 2004 diabetic foot infections guidelines.

Research since the previous guidelines shows that many foot infections are treated improperly, including prescribing the wrong antibiotic or not addressing underlying conditions such as peripheral arterial disease.

The new guidelines include 10 common questions with extensive, evidence-based answers, which the panel that wrote the guidelines determined were most likely to help a health care provider treating a patient with [diabetes](#) who has a foot wound. The first step is to determine if the wound is infected, which the guidelines note is likely if there are at least two of the following signs: redness, warmth, tenderness, pain or swelling. About half of ulcers are not infected and therefore should not be treated with antibiotics, the guidelines note. People with infections do need antibiotic therapy and those with a severe infection

should be hospitalized immediately.

When a foot sore is infected, imaging the foot is usually necessary to determine if the bone is infected. It is also important to perform a culture of the wound to determine the bacteria causing the infection, which will then help guide which antibiotic should be used for treatment. Because of the complexity of diabetic [foot infections](#), the guidelines suggest these patients are best served by a multidisciplinary team, including infectious diseases specialists, podiatrists, surgeons and orthopedists. In rural areas, doctors may be able to use telemedicine to consult with the appropriate experts, Dr. Lipsky said.

"There is quite a bit of over-prescribing or inappropriate prescribing of antibiotics for diabetic foot wounds, which doesn't help the patient and can lead to antibiotic resistance," said Warren S. Joseph, DPM, co-author of the guidelines and consultant for lower extremity [infectious diseases](#) at Roxborough Memorial Hospital, Philadelphia. "The guidelines note that when antibiotics are necessary they should be discontinued when the infection is gone, even if the wound hasn't completely healed."

The voluntary [guidelines](#) are not intended to take the place of a doctor's judgment, but rather to support the decision-making process, which must be individualized according to each patient's circumstances.

Provided by Infectious Diseases Society of America

Citation: Correct treatment of common diabetic foot infections can reduce amputations (2012, May 22) retrieved 6 May 2024 from <https://medicalxpress.com/news/2012-05-treatment-common-diabetic-foot-infections.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.