


BMI thresholds for gestational diabetes differ by race

June 1 2012



There is considerable racial/ethnic variation in the prevalence of gestational diabetes mellitus by body mass index, according to a study published online May 22 in *Diabetes Care*.

(HealthDay) -- There is considerable racial/ethnic variation in the prevalence of gestational diabetes mellitus (GDM) by body mass index (BMI), according to a study published online May 22 in *Diabetes Care*.

Monique Hedderson, Ph.D., from Kaiser Permanente Northern California in Oakland, and colleagues analyzed data from 123,040 women without diabetes prior to their pregnancy, who delivered babies between 1995 and 2006. The racial/[ethnic disparities](#) in the prevalence of GDM by BMI were assessed.

The researchers found that the age-adjusted prevalence of GDM increased with increasing BMI in all racial/ethnic groups. The prevalence

of GDM differed by race; at a BMI of 22.0 to 24.9 kg/m², the prevalence was 9.9 percent for Asian women and 8.5 percent for Filipina women, whereas for Hispanic, non-Hispanic white, and African-American women, the prevalence of GDM was >8.0 percent at a BMI of 28 to 30, 34 to 36, and ≥37 kg/m², respectively. If all pregnant women were of normal weight (BMI

Citation: BMI thresholds for gestational diabetes differ by race (2012, June 1) retrieved 24 April 2024 from <https://medicalxpress.com/news/2012-06-bmi-thresholds-gestational-diabetes-differ.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.