

# MRI scans show how sleep loss affects the ability to choose proper foods

June 10 2012

---

MRI scans from a study being presented today at SLEEP 2012 reveal how sleep deprivation impairs the higher-order regions in the human brain where food choices are made, possibly helping explain the link between sleep loss and obesity that previous research has uncovered.

Twenty-three healthy adults participated in two sessions using [functional magnetic resonance imaging](#) (fMRI), one after a normal night's sleep and a second after a night of [sleep deprivation](#). In both sessions, participants rated how much they wanted various food items shown to them while they were inside the scanner.

"Our goal was to see if specific regions of the brain associated with food processing were disrupted by sleep deprivation," said lead author Stephanie Greer, a graduate student at the Sleep and Neuroimaging Laboratory at the University of California, Berkeley.

Results show that sleep deprivation significantly impaired brain activity in the frontal lobe, a region critical for controlling behavior and making complex choices, such as the selection of food to eat. The study suggests that sleep loss may prevent the higher brain functions normally critical for making appropriate [food choices](#), rather than necessarily changing activity in deeper brain structures that react to basic desire.

"We did not find significant differences following sleep deprivation in [brain areas](#) traditionally associated with basic reward reactivity," Greer said. "Instead, it seems to be about the regions higher up in the brain,

specifically within the frontal lobe, failing to integrate all the different signals that help us normally make wise choices about what we should eat."

She added that this failure of the frontal lobe to optimally gather the information needed to decide on the right types of foods to eat – such as how healthy relative to how tasty an item may be – may represent one brain mechanism explaining the link between [sleep loss](#) and obesity.

"These results shed light on how the brain becomes impaired by sleep deprivation, leading to improper food choices," Greer said.

**More information:** The abstract "Sleep deprivation disrupts human brain reactivity in response to food desire" is being presented today at SLEEP 2012, the 26th annual meeting of the Associated Professional Sleep Societies (APSS) in Boston.

Provided by American Academy of Sleep Medicine

Citation: MRI scans show how sleep loss affects the ability to choose proper foods (2012, June 10) retrieved 27 April 2024 from <https://medicalxpress.com/news/2012-06-mri-scans-loss-affects-ability.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------