

Post-stroke depression linked to functional brain impairment

June 5 2012

Researchers studying stroke patients have found a strong association between impairments in a network of the brain involved in emotional regulation and the severity of post-stroke depression. Results of the study are published online in the journal *Radiology*.

"A third of [patients](#) surviving a stroke experience post-stroke depression (PSD)," said lead researcher Igor Sibon, M.D., Ph.D., professor of neurology at the University of Bordeaux in Bordeaux, France. "However, studies have failed to identify a link between lesions in the [brain](#) caused by ischemia during a stroke and subsequent depression."

Instead of looking for dysfunction in a specific area of the brain following a stroke, Dr. Sibon's study was designed to assess a group of brain structures organized in a functional network called the default-mode network (DMN). Modifications of connectivity in the DMN, which is associated with internally generated thought processes, has been observed in depressive patients.

"The default-mode network is activated when the brain is at rest," Dr. Sibon said. "When the brain is not actively involved in a task, this area of the brain is engaged in internal thoughts involving self-related [memory retrieval](#) and processing."

In the study, 24 patients between the ages of 18 and 80 underwent resting-state [functional magnetic resonance imaging](#) (fMRI) 10 days after having mild to moderate [ischemic stroke](#). An fMRI imaging study

measures metabolic changes in specific areas of the brain. Although many fMRI exams are designed to measure [brain changes](#) while a patient performs a specific task, during a resting-state fMRI exam, patients lie motionless.

The patients, which included 19 men and five women, were also clinically evaluated 10 days and three months post-stroke to determine the presence and severity of depression and [anxiety symptoms](#). At three months post-stroke, patients were evaluated for depression using the DSM-IV diagnostic classification system.

Using the DSM-IV criteria, 10 patients had minor to moderate depression, and 14 patients had no depression. Results of the fMRI exams revealed an association between modifications of connectivity in the DMN 10 days after stroke and the severity of depression three months post-stroke.

"We found a strong association between early resting-state network modifications and the risk of post-stroke mood disorders," Dr. Sibon said. "These results support the theory that functional brain impairment following a stroke may be more critical than structural lesions."

According to Dr. Sibon, the widespread chemical changes that result from a stroke may lead to the modification of connectivity in brain networks such as the DMN. He said results of his study may contribute to the clinical management of [stroke patients](#) by providing an opportunity to investigate the effects of a variety of treatments on patients whose fMRI results immediately post-stroke indicate impaired connectivity in the DMN.

More information: "Subacute Default Mode Network Dysfunction in the Prediction of Post-Stroke Depression Severity," *Radiology*.

Provided by Radiological Society of North America

Citation: Post-stroke depression linked to functional brain impairment (2012, June 5) retrieved 5 May 2024 from <https://medicalxpress.com/news/2012-06-post-stroke-depression-linked-functional-brain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.